

On the topology of real algebraic stacks

Emiliano Ambrosi and Olivier de Gaay Fortman

April 3, 2025

Abstract

Motivated by questions arising in the theory of moduli spaces in real algebraic geometry, we develop a range of methods to study the topology of the real locus of a Deligne–Mumford stack over the real numbers. As an application, we verify in several cases the Smith–Thom type inequality for stacks that we conjectured in an earlier work. This requires combining techniques from group theory, algebraic geometry, and topology.

1 Introduction

This work is the second in a two-part series dedicated to the study of the topology of real Deligne–Mumford stacks. In the first paper [AGF25], we established several foundational topological properties of such stacks and formulated a conjecture on their cohomology, extending the classical Smith–Thom inequality to the stack-theoretic setting. In the present paper, we develop techniques for computing the topology of various classes of real Deligne–Mumford stacks – such as finite quotient stacks and gerbes over real varieties – and apply these methods to verify the conjecture in a range of cases.

1.1 Smith–Thom inequality for real algebraic varieties. We begin by recalling the conjecture and our motivation behind it. Let X be a real algebraic variety, by which we mean a reduced and separated scheme of finite type over \mathbb{R} . One of the foundational results in real algebraic geometry (see [Flo52; Bor60; Tho65; DIK00; Man17] for various proofs) is the Smith–Thom inequality

$$h^*(X(\mathbb{R})) = \dim H^*(X(\mathbb{R}), \mathbb{Z}/2) \leq \dim H^*(X(\mathbb{C}), \mathbb{Z}/2) = h^*(X(\mathbb{C})). \quad (1)$$

It allows one to bound the cohomology of $X(\mathbb{R})$ in terms of the one of $X(\mathbb{C})$, usually much easier to compute. Here, and in the sequel, $h^*(Y)$ denotes the dimension of the cohomology ring $H^*(Y, \mathbb{Z}/2) = \oplus_{i \geq 0} H^i(Y, \mathbb{Z}/2)$ of a topological space Y .

1.2 Failure of the naive Smith–Thom inequality for real algebraic stacks. In recent years, there has been increasing interest in moduli problems over \mathbb{R} , particularly in determining whether (1) attains equality for the associated coarse moduli space. Notable cases include moduli spaces of stable vector bundles on a curve [BS22], Hilbert schemes of points on a surface [Fu23; KR24], and symmetric powers of varieties [BD17; Fra18].

Note, however, that such a study says something about the *real moduli space* associated to the moduli problem only if this real moduli space arises as the real locus of the coarse moduli space, a phenomenon which in fact seems rare. For instance, if \mathcal{A}_1 is the coarse moduli space of elliptic curves, then $\mathcal{A}_1(\mathbb{R}) = \mathbb{R}$ parametrizes complex elliptic curves that admit a real structure up to complex isomorphism, whereas the real moduli space of real elliptic curves has two connected components (there are exactly two topological types of real elliptic curves).

To bypass this limitation, and start a systematic approach to study the topology of real moduli spaces, one is led to consider real algebraic stacks. If \mathcal{X} is such a stack, then $\mathcal{X}(\mathbb{R})$ is a category rather than a set. To obtain a topological space in a way that generalizes the euclidean topology on $X(\mathbb{R})$ when X is a real variety, one considers the set $|\mathcal{X}(\mathbb{R})|$ of isomorphism classes of $\mathcal{X}(\mathbb{R})$, and defines a natural topology on $|\mathcal{X}(\mathbb{R})|$ as in [GF22b]. A similar procedure defines a topology on the set $|\mathcal{X}(\mathbb{C})|$ of isomorphism classes of $\mathcal{X}(\mathbb{C})$ (if \mathcal{X} is separated Deligne–Mumford, the latter coincides with the topology on $|\mathcal{X}(\mathbb{C})|$ induced by the coarse moduli space).

The advantage of this perspective is that when the algebraic stack \mathcal{X} represents a moduli problem—parametrizing equivalence classes of certain algebraic objects (such as genus g curves or sheaves on a fixed variety)—the set $|\mathcal{X}(\mathbb{R})|$ corresponds to the real isomorphism classes of the real objects. For instance, $|\mathcal{M}_g(\mathbb{R})|$ represents the space of isomorphism classes of real algebraic curves of genus g .

In order to understand the topological properties of $|\mathcal{X}(\mathbb{R})|$, a natural first step is to understand whether the foundational inequality (1) generalizes to this setting. As already mentioned in [AGF25], this is not the case, as the moduli space \mathcal{A}_1 of elliptic curves example shows: we have $h^*(|\mathcal{A}_1(\mathbb{C})|) = 1$ while $h^*(|\mathcal{A}_1(\mathbb{R})|) = 2$.

The main challenge in extending the Smith–Thom inequality (1) to algebraic stacks is that, although $|\mathcal{X}(\mathbb{C})|$ is equipped with an involution $\sigma: |\mathcal{X}(\mathbb{C})| \rightarrow |\mathcal{X}(\mathbb{C})|$ which

generalizes complex conjugation on the complex locus of a real variety, the natural map

$$|\mathcal{X}(\mathbb{R})| \rightarrow |\mathcal{X}(\mathbb{C})|^\sigma \quad (2)$$

is, even in simple cases, neither injective (Example 3.3) nor surjective (Example 3.4). This can be explained as follows.

- The failure of surjectivity of (2) is due to the existence of isomorphism classes of objects $x \in \mathcal{X}(\mathbb{C})$ which are isomorphic to their complex conjugate, but not defined over \mathbb{R} .
- The failure of injectivity is measured by the following observation: for $x \in \mathcal{X}(\mathbb{R})$, the fiber of (2) above the image of x in $|\mathcal{X}(\mathbb{C})|^\sigma$ is in canonical bijection with the first non-abelian Galois cohomology group $H^1(G, \text{Aut}(x_{\mathbb{C}}))$, where

$$G := \text{Gal}(\mathbb{C}/\mathbb{R}) \cong \mathbb{Z}/2.$$

In a sense, the topological space $|\mathcal{X}(\mathbb{C})|$ is therefore too small to fully encode information about $|\mathcal{X}(\mathbb{R})|$, as it does not capture, for instance, the automorphisms of objects in $\mathcal{X}(\mathbb{C})$. To take these into account, we considered the inertia stack $\mathcal{I}_{\mathcal{X}}$, whose complex locus $\mathcal{I}_{\mathcal{X}}(\mathbb{C})$ consists of pairs (x, ϕ) with $x \in \mathcal{X}(\mathbb{C})$ and ϕ an automorphism of x . With the above considerations in mind, we proposed in [AGF25] the following conjectural generalization of the Smith–Thom inequality (1) to real Deligne–Mumford stacks.

Conjecture 1.1 (cf. [AGF25], Conjecture 1.7). *Let \mathcal{X} be a separated Deligne–Mumford stack of finite type over \mathbb{R} , with inertia $\mathcal{I}_{\mathcal{X}} \rightarrow \mathcal{X}$. Then the following inequality holds:*

$$\dim H^*(|\mathcal{X}(\mathbb{R})|, \mathbb{Z}/2) \leq \dim H^*(|\mathcal{I}_{\mathcal{X}}(\mathbb{C})|, \mathbb{Z}/2). \quad (3)$$

Recall that if \mathcal{X} is a scheme, then $\mathcal{I}_{\mathcal{X}} \rightarrow \mathcal{X}$ is an isomorphism. Thus, in that case, (3) gives back the classical Smith–Thom inequality (1).

We warn the reader that, in general, there is no natural embedding of $|\mathcal{X}(\mathbb{R})|$ into $|\mathcal{I}_{\mathcal{X}}(\mathbb{C})|$. For example, take an elliptic curve E over \mathbb{R} such that $h^*(E(\mathbb{R})) = 4$, and consider the stacky quotient $\mathcal{X} := [E/\langle -1 \rangle]$, where $-1: E \rightarrow E$ is the multiplication by -1 . One can show (see Example 3.6) that $|\mathcal{X}(\mathbb{R})| \simeq E(\mathbb{R}) \amalg E(\mathbb{R})$, and that $|\mathcal{I}_{\mathcal{X}}(\mathbb{C})| \simeq E(\mathbb{C}) \amalg \left(\coprod_{x \in E(\mathbb{C})[2]} \{x\} \right)$. The situation is depicted in Figure 1 below. Although there is no natural embedding $E(\mathbb{R}) \amalg E(\mathbb{R}) \rightarrow |\mathcal{I}_{[E/\langle -1 \rangle]}(\mathbb{C})|$, inequality (3) holds in this case (and is an equality): we have $h^*(|\mathcal{X}(\mathbb{R})|) = 8$ and $h^*(|\mathcal{I}_{\mathcal{X}}(\mathbb{C})|) = 4 + 4 = 8$.




Figure 1: The real locus and inertia of the stack $[E/\langle -1 \rangle]$.

1.3 Topology of real quotient stacks. A distinctive feature of the classical Smith–Thom inequality (1) is its inherently global nature: since varieties are locally contractible, the inequality holds trivially at the local level. In contrast, the inequality proposed in Conjecture 1.1 does not seem to be locally trivial. Indeed, any separated Deligne–Mumford stack over \mathbb{R} is étale locally isomorphic to a quotient stack $[X/\Gamma]$, where X is a real algebraic variety and Γ is a finite group scheme Γ over \mathbb{R} acting on X over \mathbb{R} , but even for $[X/\Gamma]$, Conjecture 1.1 does not appear to be straightforward. In fact, as it turns out, the topology of a real quotient stack can be quite complicated as the following theorem shows.

Let Γ be a finite group scheme over \mathbb{R} , so that G acts on $\Gamma(\mathbb{C})$ via an involution $\sigma: \Gamma(\mathbb{C}) \rightarrow \Gamma(\mathbb{C})$, and let X be a quasi-projective scheme over \mathbb{R} on which Γ acts. Choose a complete set $H \subset \Gamma(\mathbb{C})$ of representative for the non-abelian Galois cohomology group

$$H^1(G, \Gamma(\mathbb{C})) = \{\gamma \in \Gamma \text{ such that } \gamma\sigma(\gamma) = e\} / \sim$$

where \sim is equivalence relation $\gamma \sim \beta\gamma\sigma(\beta)^{-1}$ for $\beta \in \Gamma(\mathbb{C})$. Via a twisting procedure, to each $\gamma \in H$ one can associate a finite group scheme Γ_γ over \mathbb{R} , a quasi-projective scheme X_γ over \mathbb{R} , and a natural action of the group scheme Γ_γ on the quasi-projective scheme X_γ over \mathbb{R} (see Section 3.1.2 for details).

The following is the first main result of the paper.

Theorem 1.2. *Consider the above notation. There is a canonical homeomorphism*

$$|[X/\Gamma](\mathbb{R})| \xrightarrow{\sim} \coprod_{\gamma \in H} X_\gamma(\mathbb{R})/\Gamma_\gamma(\mathbb{R}).$$

We use Theorem 1.2 to prove Conjecture 1.1 in a number of examples. The following corollary (Examples 3.6 & 3.7) yields non-trivial cases of stacks of arbitrary dimension,

which are not products of lower dimensional stacks, for which Conjecture 1.1 holds.

Corollary 1.3. *Let \mathcal{X} be a Deligne–Mumford stack over \mathbb{R} . Assume that one of the following two conditions holds:*

1. *We have $\mathcal{X} = [(X \times X)/\mathbb{Z}/2]$ for a real variety X , where $\mathbb{Z}/2$ acts on $X \times X$ by permuting the factors.*
2. *We have $\mathcal{X} = [A/\langle -1 \rangle]$, where A is an abelian variety over \mathbb{R} and $-1: A \rightarrow A$ the multiplication by -1 homomorphism.*

Then Conjecture 1.1 holds for \mathcal{X} .

1.4 Positive results for stacky curves. With Theorem 1.2 in our hands, we then move to prove the Conjecture 1.1 for a large class of stacky curves. This is the content of the second main result of the paper. To state this result, we define a *real curve* as a one-dimensional variety over \mathbb{R} (see Section 2), not necessarily proper.

Theorem 1.4. *Let X be a smooth real curve, and let Γ be a finite group scheme over \mathbb{R} which acts on X over \mathbb{R} . Assume that one of the following conditions holds:*

1. *The action of Γ on X is faithful.*
2. *The group scheme Γ is abelian.*

Then Conjecture 1.1 holds for the quotient stack $\mathcal{X} = [X/\Gamma]$.

The proof of Theorem 1.4 is somewhat indirect, as we do not directly compare the topological spaces $|[X/\Gamma](\mathbb{R})|$ and $|\mathcal{I}_{[X/\Gamma]}(\mathbb{C})|$. Instead, we compute $h^*([X/\Gamma](\mathbb{R}))$ and $h^*(|\mathcal{I}_{[X/\Gamma]}(\mathbb{C})|)$ separately by combining local and global methods. One of the main ingredients in the argument is the study of the local structure of the natural morphism $|[X/\Gamma](\mathbb{R})| \rightarrow (X/\Gamma)(\mathbb{R})$, which we use to analyze the global geometry of $|[X/\Gamma](\mathbb{R})|$. We then compare the two numbers $h^*([X/\Gamma](\mathbb{R}))$ and $h^*(|\mathcal{I}_{[X/\Gamma]}(\mathbb{C})|)$ using the classical Smith–Thom inequality (1) and a group-theoretic inequality from [AGF25].

1.5 Topology of split gerbes over a real variety. Finally, we try to go beyond the case of quotients of a real variety by a finite group scheme over \mathbb{R} , by considering split gerbes over a real variety. These are stacks of the form $\mathcal{X} = [U/H]$, where U is a real variety and $H \rightarrow U$ a finite étale group scheme over U (and where the action of H on U over U is the trivial action). Split gerbes over a real variety seem important in the study of the topology of real Deligne–Mumford stacks in general, and of Conjecture 1.1 in particular, as any Deligne–Mumford stack \mathcal{X} over \mathbb{R} admits a stratification whose strata are étale locally of the form $[U/H]$.

In the following theorem, which is the third and last main result of this paper, we describe the topology of the real locus of such a stack.

Theorem 1.5. *Let U be a geometrically connected \mathbb{R} -variety with $U(\mathbb{R}) \neq \emptyset$. Let $H \rightarrow U$ be a finite étale group scheme. For $p \in U(\mathbb{C})$, consider the induced action of the fundamental group $\pi_1(U(\mathbb{C}), p)$ on the fiber $H_p(\mathbb{C})$. The following holds.*

1. *The canonical map $f: |[U/H](\mathbb{R})| \rightarrow U(\mathbb{R})$ is a topological covering over each connected component of $U(\mathbb{R})$, with fiber $H^1(G, H_p(\mathbb{C}))$ above a point $p \in U(\mathbb{R})$.*
2. *Let C be a connected component of $U(\mathbb{R})$, and fix $p \in C$. The image of the natural map $\pi_1(C, p) \rightarrow \pi_1(U(\mathbb{C}), p)$ lies in the subgroup of elements $g \in \pi_1(U(\mathbb{C}), p)$ whose action on $H_p(\mathbb{C})$ is G -equivariant. In particular, the group $\pi_1(C, p)$ acts naturally on $H^1(G, H_p(\mathbb{C}))$.*
3. *The covering space associated to the above action of $\pi_1(C, p)$ on $H^1(G, H_p(\mathbb{C}))$ is canonically isomorphic to the covering space $f^{-1}(C) \rightarrow C$.*

To prove Theorem 1.5, we analyze the relationship between the action on the fibers of the morphism $H|_{U(\mathbb{R})} \rightarrow U(\mathbb{R})$ by the topological fundamental groups $\pi_1(C_i, p_i)$ of the connected components C_i of $U(\mathbb{R})$ and the étale fundamental group $\pi_1^{\text{ét}}(U, \bar{p}_i)$ of U . In particular, we show that the knowledge of the action of G on $U(\mathbb{C})$ and on $H_p(\mathbb{C})$ for a single base point $p \in U(\mathbb{R})$ allows one to compute the action of G on $H_q(\mathbb{C})$ for any other point $q \in U(\mathbb{R})$, see Proposition 5.3. This result may be of independent interest.

In Section 6, we reformulate the abstract statement of Theorem 1.5 in more concrete group-theoretic terms; see Proposition 6.2. We then explicitly compute the topological covering appearing in item 3 of Theorem 1.5 in several examples. This allows us to prove the Smith–Thom inequality (1) for various gerbes over an Enriques surface.

As a further application of Theorem 1.5, we establish the Smith–Thom inequality for a certain class of split gerbes over a real curve.

Corollary 1.6. *Let U be smooth curve over \mathbb{R} and $n \in \mathbb{Z}_{\geq 0}$. Let $H \rightarrow U$ be a finite étale group scheme with geometric fibers isomorphic to $(\mathbb{Z}/2)^n$. Then Conjecture 1.1 holds for the stack $[U/H]$.*

Moreover, in the proper case, Theorem 1.5 yields the following result.

Corollary 1.7. *Let U be smooth proper curve over \mathbb{R} of genus g and fix $p \in U(\mathbb{R})$. Let $H \rightarrow U$ be a finite étale group scheme. Let C_1, \dots, C_m be the connected components of $U(\mathbb{R})$ and for each i choose a point $p_i \in C_i$. Consider the natural action of $\pi_1(C_i, p_i)$*

on $H^1(G, H_{p_i}(\mathbb{C}))$, see Theorem 1.5. If Conjecture 1.1 holds for the stack $[U/H]$, then

$$\sum_{i=1}^m \# \left(\frac{H^1(G, H_{p_i}(\mathbb{C}))}{\pi_1(C_i, p_i)} \right) \leq 2 \cdot \# \left(\frac{H_p(\mathbb{C})}{\pi_1(U(\mathbb{C}), p)} \right) + (g-1) \cdot \# H_p(\mathbb{C}). \quad (4)$$

Observe that if $H \rightarrow U$ is the identity, the inequality (4) is the classical Smith-Thom inequality. By Corollary 1.6, inequality (4) holds when the fibers of $H \rightarrow U$ are of the form $(\mathbb{Z}/2)^n$ for some $n \geq 0$. We do not know a direct proof of this inequality without passing through the geometry of $[U/H]$, i.e., without using Corollary 1.6.

1.6 Organization of the paper. This paper is organized as follows. In Section 2, we fix the conventions and notation used throughout the paper. In Section 3, we prove a formula for the real locus of a quotient stack, which we then apply to verify Conjecture 1.1 in numerous examples. In Section 4, we prove the conjecture for a broad class of real stacky curves. Section 5 focuses on the topology of split gerbes, and we use this analysis to establish the conjecture for various split gerbes over a real curve. Finally, in Section 6, we reformulate the results of Section 5 in group-theoretic terms, and then use this to prove Conjecture 1.1 for several split gerbes over an Enriques surface.

1.7 Acknowledgements. We thank Olivier Benoist, Matilde Manzaroli and Florent Schaffhauser for helpful discussions. Special thanks to Matilde Manzaroli for helping us with the pictures in the paper. This research was partly supported by the grant ANR-23-CE40-0011 of Agence National de la Recherche. The second author has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme under grant agreement N°948066 (ERC-StG RationAlgic), and from the ERC Consolidator Grant FourSurf N°101087365.

2 Notation and preliminaries

2.1 Sets and topological spaces. If X is a finite set, we write $\#X$ for the cardinality of X . All the topological spaces in this paper are assumed to be locally compact and Hausdorff. If X is a topological space such that $\dim H^*(X, \mathbb{Z}/2)$ is finite, we define $h^*(X) = \dim H^*(X, \mathbb{Z}/2)$.

A *circle* is a topological space homeomorphic to $\mathbb{S}^1 := \{z \in \mathbb{C} \mid |z| = 1\}$. An *interval* is a topological space homeomorphic to one of intervals $(0, 1)$, $[0, 1)$ or $[0, 1]$ inside \mathbb{R} . If the topological space is homeomorphic to $(0, 1)$ we call it *open interval*.

2.2 Groups. Throughout the paper, G will be the finite group $G = \text{Gal}(\mathbb{C}/\mathbb{R}) \cong \mathbb{Z}/2$, and $\sigma \in G$ denotes a generator. Let Γ be a group on which G -acts. Recall (see e.g. [Ser94, Chapitre I, §5]) that the non-abelian Galois cohomology group $H^1(G, \Gamma)$ can be canonically identified with $Z^1(G, \Gamma)/\sim$ where $Z^1(G, \Gamma) \subset \Gamma$ is the set of $\gamma \in \Gamma$ such that $\gamma\sigma(\gamma) = e$ and \sim is the equivalence relation $\gamma \sim \beta\gamma\sigma(\beta)^{-1}$ for $\beta \in \Gamma(\mathbb{C})$.

If X is a set on which a group H acts, we write X/H for the set of orbits. Unless stated otherwise, we consider H acting on itself by conjugation, so that H/H is the set of conjugacy classes of H .

2.3 Real varieties. A *variety* over \mathbb{R} (resp. \mathbb{C}) will be a reduced and separated scheme of finite type over \mathbb{R} (resp. \mathbb{C}). A variety over \mathbb{R} will also be called a *real variety*. If X is a real variety, we denote by $\sigma_X: X(\mathbb{C}) \rightarrow X(\mathbb{C})$ the induced anti-holomorphic involution, which we also view as an action of G on $X(\mathbb{C})$.

A *curve* over \mathbb{R} (resp. \mathbb{C}) is a variety of dimension one over \mathbb{R} (resp. \mathbb{C}). Note that we do not assume that X is proper. For a smooth curve X over \mathbb{R} , any connected component $C \subset X(\mathbb{R})$ is homeomorphic to either the circle or the open interval.

For $n \in \mathbb{Z}_{\geq 1}$, we let μ_n be the \mathbb{R} -group scheme with $\mu_n(S) = \{x \in \mathcal{O}_S(S) \mid x^n = 1\}$ for a scheme S over \mathbb{R} .

2.4 Algebraic stacks. We indicate an algebraic stack by a calligraphic letter, such as $\mathcal{X}, \mathcal{Y}, \mathcal{Z}$. Schemes are usually indicated by roman capitals, such as X, Y, Z . For an algebraic stack \mathcal{X} , we let $\mathcal{I}_{\mathcal{X}} \rightarrow \mathcal{X}$ denote the inertia stack over \mathcal{X} .

When \mathcal{X} is an algebraic stack over a scheme S , we let $|\mathcal{X}(S)|$ denote the set of isomorphism classes of the groupoid $\mathcal{X}(S)$. For an algebraic stack \mathcal{X} over \mathbb{R} , and an object $x \in \mathcal{X}(\mathbb{R})$, let $x_{\mathbb{C}} \in \mathcal{X}(\mathbb{C})$ denote the pull-back of x along $\text{Spec}(\mathbb{C}) \rightarrow \text{Spec}(\mathbb{R})$.

Let \mathcal{X} be a Deligne-Mumford stack over \mathbb{R} . Recall that the sets $|\mathcal{X}(\mathbb{R})|$ and $|\mathcal{X}(\mathbb{C})|$ are endowed with a natural topology. To define it, one chooses a scheme U and an étale morphism $U \rightarrow \mathcal{X}$ such that

$$U(\mathbb{R}) \rightarrow |\mathcal{X}(\mathbb{R})| \quad \text{and} \quad U(\mathbb{C}) \rightarrow |\mathcal{X}(\mathbb{C})|$$

are surjective. Then the topologies on $|\mathcal{X}(\mathbb{R})|$ and $|\mathcal{X}(\mathbb{C})|$ are the induced quotient topologies. See [GF22a, Section 2.3] for more details.

2.5 Preliminaries from [AGF25]. For the convenience of the reader, we recall the following results from [AGF25], that will be used several times in the sequel.

Proposition 2.1. [AGF25, Proposition 5.5]. *Let \mathcal{X} be a separated Deligne–Mumford*

stack of finite type over \mathbb{R} , with coarse moduli space $p: \mathcal{X} \rightarrow M$.

1. Let $f: |\mathcal{X}(\mathbb{R})| \rightarrow M(\mathbb{R})$ denote the map induced by p . Let $x \in \mathcal{X}(\mathbb{R})$ with isomorphism class $[x] \in |\mathcal{X}(\mathbb{R})|$. There is a canonical bijection $f^{-1}(f([x])) = H^1(G, \text{Aut}(x_{\mathbb{C}}))$.
2. We have $\#H^1(G, \text{Aut}(x_{\mathbb{C}})) = \#H^1(G, \text{Aut}(x'_{\mathbb{C}}))$ for each pair of objects $x, x' \in \mathcal{X}(\mathbb{R})$ whose induced objects $x_{\mathbb{C}}, x'_{\mathbb{C}} \in \mathcal{X}(\mathbb{C})$ are isomorphic in $\mathcal{X}(\mathbb{C})$.

Proposition 2.2. [AGF25, Proposition 5.4]. Let S be a complex variety. Let $f: X \rightarrow S$ be a scheme of finite type over S . Let $H \rightarrow S$ be a finite group scheme over S , acting on X over S . Then

$$|\mathcal{I}_{[X/H]}(\mathbb{C})| = \left\{ (x, \gamma) \in X(\mathbb{C}) \times H(\mathbb{C}) \mid \gamma \in \text{Stab}_{H_{f(x)}(\mathbb{C})}(x) \right\} / \sim \quad (5)$$

where $(x, \gamma) \sim (gx, g\gamma g^{-1})$ for $g \in H_{f(x)}(\mathbb{C})$.

3 Topology of a real quotient stack

In this section, we describe the topology of the real locus of the quotient stack $[X/\Gamma]$ of a real variety X on which a finite \mathbb{R} -group scheme Γ acts, and prove Theorem 1.2. We then use this description to verify the Smith–Thom inequality (3) in many examples.

3.1 The real locus of a quotient stack. The goal of Section 3.1 is to prove Theorem 1.2.

3.1.1 Group schemes over the reals and torsors. Let Γ be a finite group scheme over \mathbb{R} and let $\sigma_{\Gamma}: \Gamma(\mathbb{C}) \rightarrow \Gamma(\mathbb{C})$ be the action of G on $\Gamma(\mathbb{C})$ corresponding to Γ .

Choose a complete set of representative $H \subset Z^1(G, \Gamma)$ for $H^1(G, \Gamma)$, so that the composition $H \subset Z^1(G, \Gamma) \rightarrow H^1(G, \Gamma)$ is a bijection; we choose H such that $e \in H$. For each $\gamma \in H$, we define an involution

$$\varphi^{\gamma}: \Gamma(\mathbb{C}) \rightarrow \Gamma(\mathbb{C}) \quad \text{as} \quad \varphi^{\gamma}(g) = \sigma(g) \cdot \gamma^{-1}.$$

We consider the resulting G -set $(\Gamma(\mathbb{C}), \varphi^{\gamma})$. Note that left multiplication defines an action of the G -group $(\Gamma(\mathbb{C}), \sigma_{\Gamma})$ on the G -set $(\Gamma(\mathbb{C}), \varphi^{\gamma})$. In particular, if P_{γ} is the \mathbb{R} -scheme associated to $(\Gamma(\mathbb{C}), \varphi^{\gamma})$, we get an action of the \mathbb{R} -group scheme Γ on the \mathbb{R} -scheme P_{γ} that turns the latter into a Γ -torsor. Recall the following classical result.

Lemma 3.1. *The following map is bijective:*

$$\begin{aligned} H^1(G, \Gamma) = H &\rightarrow \{\text{isomorphism classes of } \Gamma\text{-torsors over } \operatorname{Spec}(\mathbb{R})\}, \\ \gamma &\mapsto P_\gamma. \end{aligned}$$

Proof. See e.g. [Sko01, Example 2, page 13]. \square

3.1.2 The topology of the real locus of a quotient stack. We continue with the above notation. For $\gamma \in H$, define an involution

$$\sigma_\Gamma^\gamma: \Gamma(\mathbb{C}) \rightarrow \Gamma(\mathbb{C}) \quad \text{as} \quad \sigma_\Gamma^\gamma(g) := \gamma \sigma_\Gamma(g) \gamma^{-1}.$$

The pair $(\Gamma(\mathbb{C}), \sigma_\Gamma^\gamma)$ corresponds to a finite group scheme Γ_γ over \mathbb{R} . Let X be a quasi-projective scheme over \mathbb{R} with real structure $\sigma_X: X(\mathbb{C}) \rightarrow X(\mathbb{C})$, acted upon from the left by the finite group scheme Γ over \mathbb{R} . For $\gamma \in H$, define an involution $\sigma_X^\gamma: X(\mathbb{C}) \rightarrow X(\mathbb{C})$ as $\sigma_X^\gamma(x) = \gamma \cdot \sigma_X(x)$. The pair $(X(\mathbb{C}), \sigma_X^\gamma)$ corresponds to a quasi-projective scheme X_γ over \mathbb{R} . Note that

$$X_\gamma(\mathbb{R}) = X(\mathbb{C})^{\sigma_X^\gamma} \quad \text{and} \quad \Gamma_\gamma(\mathbb{R}) = \Gamma(\mathbb{C})^{\sigma_\Gamma^\gamma} \quad \text{for each } \gamma \in H.$$

Proof of Theorem 1.2. Recall that, by definition one has

$$|[X/\Gamma](\mathbb{R})| = \{\text{pairs } (P, f) \mid P \text{ a } \Gamma\text{-torsor and } f \text{ a } \Gamma\text{-equivariant morphism } P \rightarrow X\} / \cong$$

where $(P_1, f_1) \cong (P_2, f_2)$ if there exists an isomorphism of torsors $h: P_1 \xrightarrow{\cong} P_2$ such that $f_2 \circ h = f_1$.

We need to prove that there exists a canonical homeomorphism

$$|[X/\Gamma](\mathbb{R})| \xrightarrow{\sim} \coprod_{\gamma \in H} X_\gamma(\mathbb{R}) / \Gamma_\gamma(\mathbb{R}).$$

To prove this, we first observe that the action of $\Gamma(\mathbb{C})$ on $X(\mathbb{C})$ is compatible with the involutions σ_X^γ and σ_Γ^γ . Indeed, for $x \in X(\mathbb{C})$ and $g \in \Gamma(\mathbb{C})$, we have:

$$\sigma_X^\gamma(g \cdot x) = \gamma \cdot \sigma_X(g \cdot x) = \gamma \cdot \sigma_\Gamma(g) \cdot \sigma_X(x) = \gamma \cdot \sigma_\Gamma(g) \cdot \gamma^{-1} \cdot \gamma \cdot \sigma_X(x) = \sigma_\Gamma^\gamma(g) \cdot \sigma_X^\gamma(x).$$

Therefore, we obtain an action of the G -group $(\Gamma(\mathbb{C}), \sigma_\Gamma^\gamma)$ on the G -space $(X(\mathbb{C}), \sigma_X^\gamma)$.

In particular, the subgroup

$$\Gamma_\gamma(\mathbb{R}) = \Gamma(\mathbb{C})^{\sigma_\Gamma^\gamma} \subset \Gamma(\mathbb{C})$$

of elements fixed under σ_Γ^γ acts on the fixed space $X_\gamma(\mathbb{R}) = X(\mathbb{C})^{\sigma_X^\gamma} \subset X(\mathbb{C})$.

Fix $\gamma \in H$ and take any $x \in X_\gamma(\mathbb{R})/\Gamma_\gamma(\mathbb{R})$. Choose a $y \in X_\gamma(\mathbb{R})$ that lifts x and consider the $\Gamma(\mathbb{C})$ -equivariant morphism

$$f_y: \Gamma(\mathbb{C}) \rightarrow X(\mathbb{C}), \quad g \mapsto g \cdot y.$$

This morphism is compatible with the G -action φ^γ on $\Gamma(\mathbb{C})$ and with the G -action σ_X on $X(\mathbb{C})$, hence it gives rise to a Γ -equivariant morphism

$$f_y: P_\gamma \rightarrow X$$

of schemes over \mathbb{R} . Define

$$\alpha(x) := (P_\gamma, f_y) \in |[X/\Gamma](\mathbb{R})|.$$

We first show that α is well-defined, i.e., that α does not depend on the choice of the lift y of x . If $z \in X_\gamma(\mathbb{R})$ is another lift of x , then there exists a $g \in \Gamma_\gamma$ such that $y = g \cdot z$. Since $g \in \Gamma_\gamma(\mathbb{R}) = \Gamma(\mathbb{C})^{\sigma_\Gamma^\gamma}$, the morphism $g: P_\gamma \rightarrow P_\gamma$ sending h to hg is an isomorphism of torsors over \mathbb{R} , fitting into the commutative diagram

$$\begin{array}{ccc} P_\gamma & \xrightarrow{f_y} & X \\ \downarrow g & & \parallel \\ P_\gamma & \xrightarrow{f_z} & X. \end{array}$$

In particular, we have an equality of isomorphism classes $[(P_\gamma, f_y)] = [(P_\gamma, f_z)] \in |[X/\Gamma](\mathbb{R})|$. We conclude that we get a canonical map

$$\alpha: |[X/\Gamma](\mathbb{R})| \rightarrow \coprod_{\gamma \in H} X_\gamma(\mathbb{R})/\Gamma_\gamma(\mathbb{R}). \quad (6)$$

We observe that, by construction and Lemma 3.1, the map α is surjective. It is also injective since $\text{Aut}_\Gamma(P_\gamma) = \Gamma_\gamma(\mathbb{R})$. Thus, it remains to prove that α is a homeomorphism.

To see this, note that for each $\gamma \in H$, we have a natural morphism

$$X_\gamma \rightarrow [X/\Gamma]. \quad (7)$$

Namely, to give such a map is to give:

- (1) a $\Gamma \times_{\mathbb{R}} X_\gamma$ -torsor $P \rightarrow X_\gamma$ over X_γ , and
- (2) a $\Gamma \times_{\mathbb{R}} X_\gamma$ equivariant morphism $P \rightarrow X \times_{\mathbb{R}} X_\gamma$ of schemes over X_γ .

As for (1), we put $P = P_\gamma \times_{\mathbb{R}} X_\gamma$, which is a $\Gamma \times_{\mathbb{R}} X_\gamma$ -torsor by base-changing the Γ -torsor structure of $P_\gamma \rightarrow \text{Spec}(\mathbb{R})$ along $X_\gamma \rightarrow \text{Spec}(\mathbb{R})$. As for (2), we consider the morphism

$$P_\gamma \times_{\mathbb{R}} X_\gamma \rightarrow X \times_{\mathbb{R}} X_\gamma \quad (8)$$

defined via Galois descent by the map

$$\Gamma(\mathbb{C}) \times X(\mathbb{C}) \rightarrow X(\mathbb{C}) \times X(\mathbb{C}), \quad (g, x) \mapsto (gx, x),$$

which is indeed compatible with the anti-holomorphic involution $(g, x) \mapsto (\varphi^\gamma(g), \sigma_X^\gamma(x))$ on the left hand side and the anti-holomorphic involution $(x, y) \mapsto (\sigma_X(x), \sigma_X^\gamma(y))$ on the right hand side. Since the map (8) is $\Gamma \times_{\mathbb{R}} X_\gamma$ -equivariant, it yields the desired morphism (7).

We obtain a morphism

$$U := \coprod_{\gamma \in H} X_\gamma \rightarrow [X/\Gamma],$$

and, by the fact that the map α in (6) is a bijection (which has already been shown), the induced map

$$U(\mathbb{R}) = \coprod_{\gamma \in H} X_\gamma(\mathbb{R}) \rightarrow |[X/\Gamma](\mathbb{R})| \quad (9)$$

is surjective. By definition of the real analytic topology on $|[X/\Gamma](\mathbb{R})|$, see Section 2.4, it follows that the topology on $|[X/\Gamma](\mathbb{R})|$ is the quotient topology coming from the surjection (9) and the real analytic topology on $U(\mathbb{R}) = \coprod_{\gamma} X_\gamma(\mathbb{R})$. As the diagram

$$\begin{array}{ccc} \coprod_{\gamma \in H} X_\gamma(\mathbb{R}) & \xlongequal{\quad} & \coprod_{\gamma \in H} X_\gamma(\mathbb{R}) \\ \downarrow & & \downarrow \\ |[X/\Gamma](\mathbb{R})| & \xrightarrow{\alpha} & \coprod_{\gamma \in H} X_\gamma(\mathbb{R})/\Gamma_\gamma \end{array}$$

commutes, and as each quotient $X_\gamma(\mathbb{R})/\Gamma_\gamma$ carries the the quotient topology coming from $X_\gamma(\mathbb{R}) \rightarrow X_\gamma(\mathbb{R})/\Gamma_\gamma$, this proves that α is a homeomorphism as wanted. \square

3.2 Smith–Thom for various quotient stacks. In this section we apply Theorem 1.2 to prove the Smith–Thom inequality (3) in a number of examples.

Example 3.2. Let Γ be any finite \mathbb{R} -group scheme. Take $X = \text{Spec}(\mathbb{R})$ with the trivial action of Γ . Then Theorem 1.2 just says that $|[X/\Gamma](\mathbb{R})|$ is the disjoint union of $\#H^1(G, \Gamma)$ points, which also follows directly from the definitions and Lemma 3.1. (Note that we already verified the Smith–Thom inequality (3) for $[X/\Gamma]$ in [AGF25, Proposition 1.8].)

Example 3.3. Let $X := \mathbb{A}_{\mathbb{R}}^1$. Let $\Gamma := \mathbb{Z}/2$ endowed with the trivial G -action. We let Γ act on X via the map sending x to $-x$. To compute $\mathcal{X} := [\mathbb{A}_{\mathbb{R}}^1/\Gamma]$, we start by observing that $H^1(G, \Gamma)$ has two elements, e (the trivial element) and γ . One computes that $X(\mathbb{R})/\Gamma = \mathbb{R}/\pm 1 \simeq \mathbb{R}_{\geq 0}$ and also $X_\gamma(\mathbb{R})/\Gamma = i\mathbb{R}/\pm 1 \simeq i\mathbb{R}_{\geq 0}$. Hence, by Theorem 1.2, we have

$$|\mathcal{X}(\mathbb{R})| = \mathbb{R}_{\geq 0} \coprod i\mathbb{R}_{\geq 0}.$$

In conclusion, we find that $h^*(|\mathcal{X}(\mathbb{R})|) = 2$. On the other hand, by Proposition 2.2, one has $I_{[X/\Gamma]}(\mathbb{C}) \simeq \mathbb{C} \coprod \{0\}$. In particular $h^*(I_{[X/\Gamma]}(\mathbb{C})) = 2$, we see that the Smith–Thom inequality (3) holds and it is an actual equality. For completeness, we also describe the natural map $f: |[\mathbb{A}_{\mathbb{R}}^1/\Gamma](\mathbb{R})| \rightarrow (X/\Gamma)(\mathbb{R})$ (see Figure 2). Identifying $(X/\Gamma)(\mathbb{C})$ with \mathbb{C} via the map $z \mapsto z^2$, one sees that $(X/\Gamma)(\mathbb{R}) = \mathbb{R} \subset \mathbb{C}$. Under this identification, f induces homeomorphisms $X(\mathbb{R})/\Gamma = \mathbb{R}_{\geq 0} \xrightarrow{z \mapsto z^2} \mathbb{R}_{\geq 0}$ and $X(\mathbb{R})_\gamma/\Gamma = i\mathbb{R}_{\geq 0} \xrightarrow{z \mapsto z^2} \mathbb{R}_{\leq 0}$. Hence $\#f^{-1}(x) = 1$ for every $x \neq 0$ and $\#f^{-1}(0) = 2$, as predicted by Proposition 2.1.




Figure 2: The morphism $|[\mathbb{A}^1/(\mathbb{Z}/2)](\mathbb{R})| \rightarrow (\mathbb{A}^1/(\mathbb{Z}/2))(\mathbb{R})$

Example 3.4. Let $X := \mathbb{A}_{\mathbb{R}}^1$. Let $\Gamma := \mathbb{Z}/2 \times \mathbb{Z}/2$ endowed with the G -action exchanging the coordinates. We let Γ act on $\mathbb{A}_{\mathbb{R}}^1$ via $(a, b) * x := (-1)^{a+b}x$. To compute $|\mathcal{X}(\mathbb{R})| :=$

$[\mathbb{A}_{\mathbb{R}}^1/\Gamma]$, we start observing that $H^1(G, \Gamma) = 0$. Hence

$$|\mathcal{X}(\mathbb{R})| = X(\mathbb{R})/\Gamma(\mathbb{R}) = X(\mathbb{R}) = \mathbb{R}$$

since $\Gamma(\mathbb{R})$ acts trivially on $X(\mathbb{C})$. We find that $h^*(|\mathcal{X}(\mathbb{R})|) = 1$. On the other hand, by Proposition 2.2, one has $I_{[X/\Gamma]}(\mathbb{C}) \simeq \mathbb{A}^1 \coprod \mathbb{A}^1 \coprod \{0\} \coprod \{0\}$ so that $h^*(I_{\mathcal{X}(\mathbb{C})}) = 6$. Hence the Smith–Thom inequality (3) holds and it is a strict inequality.

For completeness, we describe the natural map $f: |[\mathbb{A}_{\mathbb{R}}^1/\Gamma](\mathbb{R})| \rightarrow (X/\Gamma)(\mathbb{R})$ (see Figure 3). As in Example 3.3, one identifies $(X/\Gamma)(\mathbb{R})$ with $\mathbb{R} \subset \mathbb{C}$. Under this identification, the map $f: \mathbb{R} \rightarrow \mathbb{R}$ becomes the absolute value map (hence not surjective), so that $\#f^{-1}(x) = 2$ for every $x > 0$, and $\#f^{-1}(0) = 1$, as predicted by Proposition 2.1.




Figure 3: The morphism $|[\mathbb{A}^1/(\mathbb{Z}/2 \times \mathbb{Z}/2)](\mathbb{R})| \rightarrow (\mathbb{A}^1/(\mathbb{Z}/2 \times \mathbb{Z}/2))(\mathbb{R})$

Examples 3.5. Let $X := \mathbb{A}_{\mathbb{R}}^2$.

1. Let $\Gamma := \mathbb{Z}/2$ endowed with the trivial G -action. We let Γ act on X via the map sending x to $-x$. To compute the real locus of $\mathcal{X} := [\mathbb{A}_{\mathbb{R}}^2/\Gamma]$, we start by observing that $H^1(G, \Gamma)$ has two elements, e (the neutral element) and γ . By Theorem 1.2,

$$|\mathcal{X}(\mathbb{R})| = X(\mathbb{R})/\Gamma(\mathbb{R}) \coprod X_{\gamma}(\mathbb{R})/\Gamma(\mathbb{R}).$$

One computes that $X(\mathbb{R})/\Gamma$ and $X_{\gamma}(\mathbb{R})$ are two half-planes, so that $h^*(|\mathcal{X}(\mathbb{R})|) = 2$. On the other hand, by Proposition 2.2, one has $|I_{[X/\Gamma]}(\mathbb{C})| \simeq \mathbb{A}^2 \coprod \mathbb{A}^1$ so that $h^*(|I_{[X/\Gamma]}(\mathbb{C})|) = 2$. Hence the Smith–Thom inequality (3) holds, and is an equality.

2. Let $\Gamma := \mathbb{Z}/2 \times \mathbb{Z}/2$ endowed with the G -action exchanging the coordinates. We let Γ act on $\mathbb{A}_{\mathbb{R}}^2$ via its G -equivariant quotient $\mathbb{Z}/2$, acting by exchange of coordinates. To compute $\mathcal{X} := [\mathbb{A}_{\mathbb{R}}^2/\Gamma]$, we start by observing that $H^1(G, \Gamma) = 0$, so that, by Theorem 1.2,

$$|\mathcal{X}(\mathbb{R})| = X(\mathbb{R})/\Gamma(\mathbb{R}) = \mathbb{R}^2$$

since $\Gamma(\mathbb{R})$ acts trivially on $X(\mathbb{C})$. We find that $h^*(|\mathcal{X}(\mathbb{R})|) = 1$. On the other hand, by Proposition 2.2, $I_{[X/\Gamma]}(\mathbb{C}) \simeq \mathbb{A}^2 \coprod \mathbb{A}^2 \coprod \mathbb{A}^1 \coprod \mathbb{A}^1$. In particular, $h^*(I_{[X/\Gamma]}(\mathbb{C})) = 4$. Hence the Smith–Thom inequality (3) holds, and is a strict inequality.

Example 3.6. Let A be a real abelian variety of dimension g , so that $A(\mathbb{R}) \simeq (\mathbb{S}^1)^g \times (\mathbb{Z}/2)^k$ for some $0 \leq k \leq g$ compatibly with the group structure. Consider the inversion $[-1] : A \rightarrow A$ and write $\Gamma := \mathbb{Z}/2$. Let $\mathcal{X} := [A/(\mathbb{Z}/2)]$ where $\mathbb{Z}/2$ acts via $[-1]$, and let γ be the unique non trivial element of $H^1(G, \Gamma)$. By Theorem 1.2,

$$|\mathcal{X}(\mathbb{R})| := A(\mathbb{R})/[-1] \coprod A_\gamma(\mathbb{R})/[-1].$$

By construction A_γ is the quadratic twist of A , hence $A(\mathbb{R}) \cong A_\gamma(\mathbb{R})$ as topological spaces. In particular, we get

$$|\mathcal{X}(\mathbb{R})| \simeq (\mathbb{S}^1)^g \times (\mathbb{Z}/2)^k \coprod (\mathbb{S}^1)^g \times (\mathbb{Z}/2)^k,$$

hence $h^*(|\mathcal{X}(\mathbb{R})|) = 2^{g+k+1}$. On the other hand, by Proposition 2.2, we have

$$\mathcal{I}_{\mathcal{X}}(\mathbb{C}) \simeq A(\mathbb{C})/[-1] \coprod \coprod_{x \in A(\mathbb{C})[2]} \{x\}.$$

Since $A(\mathbb{C})/[-1] \simeq A(\mathbb{C})$ and $\#A(\mathbb{C})[2] = 2^{2g}$, we get $h^*(\mathcal{I}_{\mathcal{X}}(\mathbb{C})) = 2^{2g} + 2^{2g} = 2^{2g+1}$. Since $k \leq g$, the inequality (3) is verified; it is an equality if and only if A is maximal.

Example 3.7. Let Y be a real algebraic variety, let $\Gamma := \mathbb{Z}/2$ act on $Y \times Y$ by exchanging the coordinates and let $\mathcal{X} := [(Y \times Y)/\Gamma]$. If γ is the non trivial element of $H^1(G, \Gamma)$, by Theorem 1.2, one has

$$|\mathcal{X}(\mathbb{R})| \simeq (Y(\mathbb{R}) \times Y(\mathbb{R}))/\Gamma \coprod (Y \times Y)_\gamma(\mathbb{R})/\Gamma \simeq (Y(\mathbb{R}) \times Y(\mathbb{R}))/\Gamma \coprod Y(\mathbb{C})/G.$$

Observe that

$$|\mathcal{X}(\mathbb{C})|^G \simeq (Y(\mathbb{R}) \times Y(\mathbb{R}))/\Gamma \coprod \coprod_{Y(\mathbb{R})} Y(\mathbb{C})/G,$$

where $i : Y(\mathbb{R}) \hookrightarrow |\mathcal{X}(\mathbb{C})|^G$ embeds diagonally in $Y(\mathbb{R}) \times Y(\mathbb{R})$ and naturally in $Y(\mathbb{C})/G$. If $f : |\mathcal{X}(\mathbb{R})| \rightarrow (Y \times Y/\Gamma)(\mathbb{C})^G$ is the natural morphism, the exact sequence of sheaves

$$0 \rightarrow \mathbb{Z}/2 \rightarrow f_*\mathbb{Z}/2 \rightarrow i_*\mathbb{Z}/2 \rightarrow 0,$$

shows that

$$h^*(|\mathcal{X}(\mathbb{R})|) \leq h^*(|\mathcal{X}(\mathbb{C})|^G) + h^*(Y(\mathbb{R})). \quad (10)$$

On the other hand, by Proposition 2.2,

$$|\mathcal{I}_{\mathcal{X}}(\mathbb{C})| \simeq |\mathcal{X}(\mathbb{C})| \coprod Y(\mathbb{C}),$$

so that by the classical Smith–Thom inequality for $|\mathcal{X}(\mathbb{C})| \coprod Y(\mathbb{C})$, we get

$$h^*(|\mathcal{X}(\mathbb{C})|^G) + h^*(Y(\mathbb{R})) \leq h^*(|\mathcal{X}(\mathbb{C})|) + h^*(Y(\mathbb{C})) = h^*(|\mathcal{I}_{\mathcal{X}}(\mathbb{C})|).$$

Combining this with (10), we get that the Smith–Thom inequality (3) is satisfied.

4 Smith–Thom for real stacky curves

The goal of this section is to prove Theorem 1.4. As previously noted, the proof is somewhat indirect: rather than comparing the topology of $|[X/\Gamma](\mathbb{R})|$ directly with that of $I_{[X/\Gamma]}(\mathbb{C})$, we compute $h^*(I_{[X/\Gamma]}(\mathbb{C}))$ in Section 4.1, and $h^*(|[X/\Gamma](\mathbb{R})|)$ in Section 4.2. We then combine these computations in Section 4.3 to establish Theorem 1.4.

4.1 Inertia of complex stacky curves. Let X be a smooth curve over \mathbb{C} , and let Γ be a finite abelian group which acts on X over \mathbb{C} . We let $K \subset \Gamma$ be the kernel of the homomorphism $\Gamma \rightarrow \text{Aut}_{\mathbb{C}}(X)$ associated to the Γ -action, and define $Q := \Gamma/K$. This gives a short exact sequence of finite abelian groups

$$0 \rightarrow K \rightarrow \Gamma \rightarrow Q \rightarrow 0.$$

The restriction of the action on X of Γ to K yields the trivial action of K on X , and the induced action of Q on X is faithful. Let

$$M := X/Q = X/\Gamma,$$

which is the coarse moduli space of both $[X/Q]$ and $[X/\Gamma]$.

Proposition 4.1. *In the above notation, assume that the subgroup $K \subset \Gamma$ is contained in the center of Γ , so that for every $x \in X(\mathbb{C})$ there is a natural inclusion $K \subset \Gamma_x/\Gamma_x$. Let $\Delta \subset M(\mathbb{C})$ be the branch locus of the quotient map $q: X(\mathbb{C}) \rightarrow M$, and choose a*

lift $y_x \in X(\mathbb{C})$ of each $x \in \Delta$. Then there is a canonical homeomorphism

$$|\mathcal{I}_{[X/\Gamma]}(\mathbb{C})| \simeq (K \times M(\mathbb{C})) \coprod_{x \in \Delta} (\Gamma_{y_x}/\Gamma_{y_x} - K)$$

that commutes with the natural projections onto $M(\mathbb{C})$.

Proof. We may assume that X is connected. Let $I_{[X/\Gamma]}$ be the coarse moduli space of the inertia stack $\mathcal{I}_{[X/\Gamma]}$ of $[X/\Gamma]$. It suffices to show that the map $I_{[X/\Gamma]} \rightarrow M$ has $\#K$ disjoint sections. Indeed, $I_{[X/\Gamma]} \rightarrow M$ is finite by [AGF25, Lemma 5.2], hence for each irreducible component $Z \subset I_{[X/\Gamma]}$ of dimension one, the restriction $Z \rightarrow M$ is a finite morphism of curves, hence an isomorphism if it admits a section; moreover, over the open subset of M where the stabilizer group is exactly K , the fibers of $I_{[X/\Gamma]} \rightarrow M$ have cardinality exactly $\#K$ by Proposition 2.2.

Let $S \subset X \times_{\mathbb{C}} \Gamma$ be the stabilizer group scheme associated to the action of Γ on X over \mathbb{C} , so that S can be described pointwise as

$$S = \{(x, g) \in X \times_{\mathbb{C}} \Gamma \mid g \cdot x = x\}.$$

Then Γ acts on S by $\gamma \cdot (x, g) = (\gamma \cdot x, \gamma g \gamma^{-1})$ for $\gamma \in \Gamma$ and $(x, g) \in S$. Moreover, we have a canonical isomorphism $\mathcal{I}_{[X/\Gamma]} = [S/\Gamma]$ (see e.g. [Alp25, Exercise 3.2.12]), so that, in particular, $I_{[X/\Gamma]} = S/\Gamma$.

Since K is contained in the center of Γ , to any $k \in K$ one can associate the following well defined section s_k of the canonical map $I_{[X/\Gamma]} \rightarrow M$:

$$s_k: M \rightarrow I_{[X/\Gamma]} = S/\Gamma, \quad [x] \mapsto [(x, k)]. \quad (11)$$

The sections s_k and $s_{k'}$ are disjoint for $k \neq k' \in K$, and so the proposition follows. \square

Proposition 4.2. *Assume that $K \subset \Gamma$ is contained in the center of Γ . Then*

$$h^*(|\mathcal{I}_{[X/\Gamma]}(\mathbb{C})|) = \#K \cdot h^*(M(\mathbb{C})) + \left(\sum_{x \in \Delta} \#(\Gamma_{y_x}/\Gamma_{y_x}) \right) - \#\Delta \cdot \#K. \quad (12)$$

If, in addition, Γ_{y_x} is abelian for each $x \in \Delta$, then

$$h^*(|\mathcal{I}_{[X/\Gamma]}(\mathbb{C})|) = \#K \cdot h^*(|\mathcal{I}_{[X/Q]}(\mathbb{C})|). \quad (13)$$

Proof. By Proposition 4.1, we have

$$h^*(|I_{[X/\Gamma]}(\mathbb{C})|) = \#K \cdot h^*(M(\mathbb{C})) + \sum_{x \in \Delta} (\#(\Gamma_{y_x}/\Gamma_{y_x}) - \#K), \quad (14)$$

and (14) implies (12). Applying (14) to the quotient stack $[X/Q]$ gives

$$h^*(|I_{[X/Q]}(\mathbb{C})|) = h^*(M(\mathbb{C})) + \sum_{x \in \Delta} (\#(Q_{y_x}/Q_{y_x}) - 1). \quad (15)$$

If Γ_{y_x} is abelian for each $x \in \Delta$, then one has

$$\Gamma_{y_x}/\Gamma_{y_x} = \Gamma_{y_x}, \quad Q_{y_x}/Q_{y_x} = Q_{y_x}, \quad \text{and} \quad \#K \cdot \#Q_{y_x} = \#\Gamma_{y_x}.$$

Therefore, (13) follows from (14) and (15), and we are done. \square

Example 4.3. Consider the moduli stack \mathcal{A}_1 of elliptic curves over \mathbb{C} , with coarse moduli space $\mathcal{A}_1 \rightarrow \mathbf{A}_1 = \mathbb{A}_{\mathbb{C}}^1$. Then $h^*(|\mathcal{I}_{\mathcal{A}_1}(\mathbb{C})|) = 8$. Indeed, we let $\ell \geq 3$ be a prime number and let $\mathcal{A}_1[\ell]$ be the moduli space of elliptic curves with level ℓ structure; it is equipped with a $\mathrm{SL}_2(\mathbb{F}_\ell)$ -action such that $\mathcal{A}_1 = [\mathrm{SL}_2(\mathbb{F}_\ell) \backslash \mathcal{A}_1[\ell]]$. In this case, we have $K = \langle -1 \cdot \mathrm{Id} \rangle \subset \mathrm{SL}_2(\mathbb{F}_\ell) = \Gamma$, and $\Gamma/K = \mathrm{PSL}_2(\mathbb{F}_\ell)$. The locus $\Delta \subset \mathbf{A}_1(\mathbb{C})$ of isomorphism classes of elliptic curves with automorphism group larger than $\{\pm 1\}$ consists of two points, with respective automorphism groups $\mathbb{Z}/4$ and $\mathbb{Z}/6$. Thus, Proposition 4.2 implies that $h^*(|\mathcal{I}_{\mathcal{A}_1}(\mathbb{C})|) = 2 + 4 + 6 - 2 \cdot 2 = 8$.

4.2 Topology of real stacky curves. In this section, we study the topology of $|\mathcal{X}(\mathbb{R})|$ when \mathcal{X} is a stacky quotient curve over \mathbb{R} .

4.2.1 Local geometry. Let \mathcal{X} be a separated Deligne–Mumford stack of finite type over \mathbb{R} . Let $f: |\mathcal{X}(\mathbb{R})| \rightarrow M(\mathbb{R})$ be the map induced by the coarse moduli space $\mathcal{X} \rightarrow M$.

Definition 4.4. For $x \in M(\mathbb{R})$, define $h^1(G, x) = \#H^1(G, \mathrm{Aut}(z_{\mathbb{C}}))$ where $z \in \mathcal{X}(\mathbb{R})$ is an object whose isomorphism class $[z] \in |\mathcal{X}(\mathbb{R})|$ lies in the inverse image $f^{-1}(x)$ of x . By Proposition 2.1, we have $h^1(G, x) = \#f^{-1}(x)$.

For example, when \mathcal{X} is the quotient $\mathcal{X} = [X/H]$ of a real variety X by finite group scheme H over \mathbb{R} , and when $x \in (X/H)(\mathbb{R})$ is the image of a point $y \in X(\mathbb{R})$ under the map $X(\mathbb{R}) \rightarrow (X/H)(\mathbb{R})$, then $h^1(G, x) = \#H^1(G, H_y(\mathbb{C}))$. This definition allows us to study the local topology of a stacky curve around a point with non-trivial stabilizer.

Lemma 4.5. *Let X be a smooth curve over \mathbb{R} . Let H be a finite \mathbb{R} -group scheme that acts faithfully on X over \mathbb{R} .*

1. For each $x \in X(\mathbb{R})$, there exists an integer $n \geq 1$ such that the stabilizer group scheme H_x is isomorphic to μ_n .
2. For $x \in (X/H)(\mathbb{R})$, the number $h^1(G, x)$ is equal to 1 (resp. 2) if n is odd (resp. even).

Proof. Let $\Gamma = H(\mathbb{C})$. Since the action of Γ is faithful, there are only finitely many points $x \in X(\mathbb{R})$ with non trivial stabilizer Γ_x . Since the statement is trivial for points with trivial stabilizer, we focus on the points x with $\Gamma_x \neq 0$. Choose an open neighbourhood U of x , stable under the actions of G and Γ , not containing any other point with non-trivial stabilizer, and G -equivariantly biholomorphic to an open disk centered in x endowed with the G -action given by complex conjugation. Since the group of biholomorphisms of the disk with one fixed point is isomorphic to \mathbb{S}^1 , we see that Γ_x is isomorphic to \mathbb{Z}/n for some positive integer n . Moreover, a generator $\gamma \in \Gamma_x$ acts as $\gamma(z) = e^{i\theta}z$ if z is a local coordinate around x . Since the G -action is compatible with the action of Γ , this forces a G -equivariant isomorphism $\Gamma_x \simeq \mu_n$, proving the first item of the lemma. By the fact that $\#H^1(G, \mu_n)$ equals 1 if n is odd and 2 if n is even, the second item follows from the first. \square

4.2.2 Global geometry. We now study the possible shapes of the connected components of the real locus of a real stacky quotient curve.

Proposition 4.6. *Let X be a smooth curve over \mathbb{R} . Let H be a finite étale group scheme over \mathbb{R} which acts on X over \mathbb{R} . Let C be a connected component of $|[X/H](\mathbb{R})|$. Then C is homeomorphic to either an interval in \mathbb{R} of the form $(0, 1)$, $(0, 1]$ or $[0, 1]$, or to the circle \mathbb{S}^1 . If X is proper then only the possibilities $[0, 1]$ and \mathbb{S}^1 can occur.*

We actually prove something slightly more general in the following Lemma 4.7. Observe that Proposition 4.6 follows from Theorem 1.2 and Lemma 4.7.

Lemma 4.7. *Let X be a smooth curve over \mathbb{R} . Let H be a finite étale group scheme over \mathbb{R} acting on X over \mathbb{R} . Each connected component of $X(\mathbb{R})/H(\mathbb{R})$ is homeomorphic to the interval $[0, 1] \subset \mathbb{R}$, the interval $(0, 1]$, the interval $(0, 1)$, or the circle $\mathbb{S}^1 \subset \mathbb{R}^2$.*

Proof. We may and do assume that H acts faithfully on X .

First, assume that X is proper, so that $X(\mathbb{R})$ is compact, and let C be a connected component of $X(\mathbb{R})$ with stabilizer $\text{Stab}_{H(\mathbb{R})}(C)$ in $H(\mathbb{R})$. We start by proving that

$$C/\text{Stab}_{H(\mathbb{R})}(C) \simeq \mathbb{S}^1 \quad \text{or} \quad C/\text{Stab}_{H(\mathbb{R})}(C) \simeq [0, 1]. \quad (16)$$

Recall that every connected Riemann surface S admits a unique complete Riemann metric g with constant curvature being negative (genus ≥ 2), zero (genus zero) or positive (genus one). Moreover, for genus ≥ 2 the group $\text{Bihol}(S)$ coincides with the group $\text{Isom}(S, g)^+$ of orientation preserving isometries of the Riemannian manifold (S, g) . In genus zero we have, for the subgroup $\text{PGL}_2(\mathbb{R}) \subset \text{PGL}_2(\mathbb{C}) = \text{Bihol}(\mathbb{P}^1(\mathbb{C}))$, that $\text{PGL}_2(\mathbb{R}) = \text{SO}_3(\mathbb{R})$ acts as isometries on $\mathbb{P}^1(\mathbb{C}) \cong S^2$. The automorphism group of any complex elliptic curve preserves its Riemannian metric.

In particular, as H acts faithfully on X , there are natural inclusions

$$\text{Stab}_{H(\mathbb{R})}(C) \subset H(\mathbb{R}) \subset \text{Isom}(X(\mathbb{C})) \subset \text{Homeo}(X(\mathbb{C})),$$

where $\text{Isom}(X(\mathbb{C}))$ is the group of isometries with respect to the Riemannian metric of $X(\mathbb{C})$. We endow C with the Riemannian metric induced by the embedding $C \subset X(\mathbb{C})$. Then C is a compact one-dimensional Riemannian manifold, and hence isometric to a circle of some length L : we have $C \cong \mathbb{R}/L\mathbb{Z}$ with the standard Riemannian metric. In particular, $\text{Isom}(C) \cong \text{O}(2)$. By the above, $\text{Stab}_{H(\mathbb{R})}(C) \subset \text{Isom}(X(\mathbb{C}))$, and hence $\text{Stab}_{H(\mathbb{R})}(C) \subset \text{Isom}(C) \cong \text{O}(2)$. So, $\text{Stab}_{H(\mathbb{R})}(C)$ is a finite subgroup of $\text{O}(2) = \text{Isom}(\mathbb{S}^1)$ with $\mathbb{S}^1 = \{z \in \mathbb{C} \mid |z| = 1\}$, hence it is generated by multiplication by some root of unity and, possibly, the standard complex conjugation on \mathbb{S}^1 . This yields (16).

Let C_1, \dots, C_n be the connected components of $X(\mathbb{R})$; each C_i is homeomorphic to \mathbb{S}^1 . Then $I := \{1, \dots, n\}$ admits a partition $I = I_1 \sqcup I_2 \sqcup \dots \sqcup I_k$ with $k \leq n$ such that the I_j are the orbits for the induced action of $H(\mathbb{R})$ on I . For each $j \in \{1, \dots, k\}$, choose an element $i_j \in I_j$. Let $H(\mathbb{R})_j = \text{Stab}_{H(\mathbb{R})}(C_{i_j})$ be the stabilizer of C_{i_j} in the group $H(\mathbb{R})$. Then

$$X(\mathbb{R})/H(\mathbb{R}) = \left(\prod_{i=1}^n C_i \right) / H(\mathbb{R}) = \prod_{j=1}^k \left(\left(\prod_{i \in I_j} C_i \right) / H(\mathbb{R}) \right) \cong \prod_{j=1}^k C_{i_j} / H(\mathbb{R})_j.$$

Thus, the lemma in the case where X is proper follows from (16).

In the general case, consider the smooth projective compactification $X \hookrightarrow Y$ of X . The action of H on X extends to an action of H on Y , and the natural map $X(\mathbb{R})/H(\mathbb{R}) \rightarrow Y(\mathbb{R})/H(\mathbb{R})$ is an open embedding whose complement is a finite set (possibly empty). By what has already been proved, each connected component of $Y(\mathbb{R})/H(\mathbb{R})$ is homeomorphic to $[0, 1]$ or \mathbb{S}^1 . By removing the points in $\Delta(\mathbb{R}) \subset Y(\mathbb{R})$, where $\Delta = Y - X$, we see that each connected component of $X(\mathbb{R})/H(\mathbb{R})$ is homeomorphic to $[0, 1]$, $(0, 1]$, $(0, 1)$ or \mathbb{S}^1 , and we are done. \square

4.2.3 Map to the coarse moduli space. Finally, we study the map from a real stacky curve to its coarse moduli space.

Let X be a smooth curve over \mathbb{R} . Let H be a finite \mathbb{R} -group scheme that acts on X over \mathbb{R} , with associated real structure $\sigma: H(\mathbb{C}) \rightarrow H(\mathbb{C})$. Let $p: [X/H] \rightarrow X/H = M$ be the coarse moduli space map, with induced map $f: |[X/H](\mathbb{R})| \rightarrow (X/H)(\mathbb{R}) = M(\mathbb{R})$.

Lemma 4.8. *Let $C \subset M(\mathbb{R})$ be a connected component. The following holds.*

1. *The map $f: |[X/H](\mathbb{R})| \rightarrow M(\mathbb{R})$ is closed.*
2. *If H acts faithfully on X , then the map $f^{-1}(C) \rightarrow C$ is surjective.*
3. *Assume that H acts faithfully on X , and that for each $m \in C$, we have $h^1(G, m) = 1$, with $h^1(G, m)$ as in Definition 4.4. Then $f^{-1}(C) \rightarrow C$ is a homeomorphism.*

Proof. To prove item 1, let $\gamma \in H(\mathbb{C})$ such that $\gamma\sigma(\gamma) = e$, and define X_γ and H_γ as in Section 3.1.2. By Theorem 1.2, it suffices to show that the natural map $X_\gamma(\mathbb{R})/H_\gamma(\mathbb{R}) \rightarrow (X/H)(\mathbb{R})$ is closed. Note that this map is induced by the natural map $X_\gamma \rightarrow X/H$, which induces an isomorphism $X_\gamma/H_\gamma \cong X/H$. The quotient map $X_\gamma \rightarrow X_\gamma/H_\gamma \cong X/H$ is finite, and hence $X_\gamma(\mathbb{R}) \rightarrow (X/H)(\mathbb{R})$ is closed. In particular, $X_\gamma(\mathbb{R})/H_\gamma(\mathbb{R}) \rightarrow (X/H)(\mathbb{R})$ is closed, proving item 1. To prove item 2, note that since the action is faithful and X is smooth, $f: |[X/H](\mathbb{R})| \rightarrow M(\mathbb{R})$ is surjective and closed; indeed, f is closed by item 1 and its image contains a dense open subset, so f is surjective. In particular, item 2 follows. Moreover, for $m \in C$, we have $\#f^{-1}(m) = h^1(G, m)$, see Proposition 2.1. Hence item 3 follows from items 1 and 2. \square

Proposition 4.9. *Assume that H acts faithfully on X over \mathbb{R} . Let $C \subset M(\mathbb{R})$ be a connected component and let $\mathcal{S} = \{x_1, \dots, x_n\} \subset C$ be the finite set of points such that $h^1(G, x_i) \neq 1$. Assume that $\mathcal{S} \neq \emptyset$.*

1. *Assume that C is an open interval, and fix a homeomorphism $\varphi: C \xrightarrow{\sim} (0, 1)$. There exists an homeomorphism*

$$\psi: f^{-1}(C) \xrightarrow{\sim} (0, y_1] \coprod [y_1, y_2] \coprod \cdots \coprod [y_{n-1}, y_n] \coprod [y_n, 1)$$

such that the following diagram commutes:

$$\begin{array}{ccc} f^{-1}(C) & \xrightarrow{\psi} & (0, y_1] \sqcup [y_1, y_2] \coprod \cdots \coprod [y_{n-1}, y_n] \coprod [y_n, 1) \\ \downarrow & & \downarrow \\ C & \xrightarrow{\varphi} & (0, 1), \end{array}$$

where the vertical arrows are the canonical ones and $y_i = \varphi(x_i)$.

2. Assume that C is a circle, and fix a homeomorphism $\varphi: C \xrightarrow{\sim} \mathbb{S}^1$ with $\varphi(x_1) = 1$. Then there exists a homeomorphism

$$\psi: f^{-1}(C) \xrightarrow{\sim} [0, \theta_1] \coprod [\theta_1, \theta_2] \coprod \cdots \coprod [\theta_{n-1}, 1]$$

such that the following diagram commutes:

$$\begin{array}{ccc} f^{-1}(C) & \xrightarrow{\psi} & [0, \theta_1] \coprod [\theta_1, \theta_2] \coprod \cdots \coprod [\theta_{n-1}, 1] \\ \downarrow & & \downarrow x \mapsto e^{2\pi i x} \\ C & \xrightarrow{\varphi} & \mathbb{S}^1, \end{array}$$

where the vertical arrows are the canonical ones and $\varphi(x_j) = e^{i\theta_j}$ for $j \neq 1$.

Proof. By Lemma 4.8, the map $f: f^{-1}(C) \rightarrow C$ is surjective. By Proposition 2.1 and Lemma 4.5, for each connected component $K \subset M(\mathbb{R})$, the map $f^{-1}(K) \rightarrow K$ is an isomorphism outside $\mathcal{S} \subset K$ and has two fibers above each point of \mathcal{S} . The proposition follows from this and from Proposition 4.6. \square

The content of Proposition 4.9 is illustrated in Figure 4 below. The picture on the left shows the case in which C is a circle: C is depicted as the internal circle and $f^{-1}(C)$ as the union of the three external arcs $[0, \theta_1]$, $[\theta_1, \theta_2]$ and $[\theta_2, 1]$. The picture on the right shows the case in which C is a open interval: C is depicted as the bottom interval and $f^{-1}(C)$ as the disjoint union of the three upper segments $(0, y_1]$, $[y_1, y_2]$ and $[y_2, 1)$.




Figure 4: The morphism $f^{-1}(C) \rightarrow C$.

4.3 Smith–Thom for real stacky curves. We are now ready to proceed with the:

Proof of Theorem 1.4. The action of H on X corresponds to a homomorphism

$$H \rightarrow \underline{\mathrm{Aut}}_{\mathbb{R}}(X), \quad (17)$$

where the latter denotes the automorphism group scheme of X over \mathbb{R} . Let $K \subset H$ be the kernel of (17), and let $Q = H/K$ be the quotient of H by K . The canonical map $Q \rightarrow \underline{\mathrm{Aut}}_{\mathbb{R}}(X)$ is a closed immersion. In particular, the group $Q(\mathbb{C})$ acts faithfully on $X(\mathbb{C})$. Let $[X/H] \rightarrow M$ be the coarse moduli space; we have $M(\mathbb{C}) = X(\mathbb{C})/H(\mathbb{C}) = X(\mathbb{C})/Q(\mathbb{C})$.

Step 1: *If H is abelian, and if the Smith–Thom inequality (3) holds for the quotient stack $[X/Q]$, then it also holds for $[X/H]$.*

Proof. Assume the Smith–Thom inequality (3) for $[X/Q]$, and consider the canonical map

$$g: |[X/H](\mathbb{R})| \rightarrow |[X/Q](\mathbb{R})|. \quad (18)$$

We claim that

$$\#g^{-1}(p) \leq \#K(\mathbb{C}) \quad \text{for any } p \in |[X/Q](\mathbb{R})|. \quad (19)$$

To prove this, choose a complete set of representatives $R_H \subset Z^1(G, H)$ for $H^1(G, H)$. Let $I_Q = m(R_H) \subset Z^1(G, Q)$ be the image of R_H in $Z^1(G, Q)$ under the natural map $m: Z^1(G, H) \rightarrow Z^1(G, Q)$, and extend I_Q to a complete set of representatives $I_Q \subset R_Q \subset Z^1(G, Q)$ for $H^1(G, Q)$. By construction, the map $m: Z^1(G, H) \rightarrow Z^1(G, Q)$ restricts to a map $m: R_H \rightarrow R_Q$. By Theorem 1.2, of which we retain the notation, we have a commutative diagram

$$\begin{array}{ccc} |[X/H](\mathbb{R})| & \xrightarrow{g} & |[X/Q](\mathbb{R})| \\ \downarrow \wr & & \downarrow \wr \\ \coprod_{\gamma \in R_H} X_{\gamma}(\mathbb{R})/H_{\gamma}(\mathbb{R}) & \longrightarrow & \coprod_{\mu \in R_Q} X_{\mu}(\mathbb{R})/Q_{\mu}(\mathbb{R}) \end{array}$$

in which the vertical arrows are homeomorphisms and where the map on the bottom is

induced by the map $m: \mathbf{R}_H \rightarrow \mathbf{R}_Q$ and the quotient morphisms

$$X_\gamma(\mathbb{R})/H_\gamma(\mathbb{R}) \rightarrow X_{m(\gamma)}(\mathbb{R})/Q_{m(\gamma)}(\mathbb{R}).$$

To prove the claim, we may assume that $p = g(q)$ for some $q \in |[X/H](\mathbb{R})|$. Let $\gamma \in \mathbf{R}_H$ such that $q \in X_\gamma(\mathbb{R})/H_\gamma(\mathbb{R})$. The exact sequence of abelian groups

$$0 \rightarrow K_\gamma(\mathbb{R}) \rightarrow H_\gamma(\mathbb{R}) \rightarrow Q_{m(\gamma)}(\mathbb{R}) \rightarrow H^1(G, K_\gamma) \rightarrow H^1(G, H_\gamma) \rightarrow H^1(G, Q_{m(\gamma)}),$$

where $K_\gamma := \text{Ker}(H_\gamma \rightarrow Q_{m(\gamma)})$, shows that $\#g^{-1}(p) \leq \#H^1(G, K_\gamma(\mathbb{C})) \leq \#K(\mathbb{C})$. This proves (19).

By Proposition 4.6, each connected component of $|[X/H](\mathbb{R})|$ or $|[X/Q](\mathbb{R})|$ is homeomorphic to a circle or an interval. Let C be a connected component of $|[X/Q](\mathbb{R})|$. If C is a circle, then, by (19), the inverse image $g^{-1}(C)$ consists of at most $\#K$ connected components which are circles or intervals. This implies that

$$h^*(g^{-1}(C)) \leq 2 \cdot \#K(\mathbb{C}) \quad \text{if } C \text{ is a circle.} \quad (20)$$

Note that every surjective continuous map from a circle to an interval has the property that the fiber above any point in the interior of the interval has cardinality at least two. To prove this, take a point p not on boundary of the interval. If p had only one pre-image, then, eliminating the point, we would get a surjective morphism from a connected space, a circle minus one point, to a non-connected one, an interval minus one point not on the boundary. This is not possible, so that p has at least two pre-images.

Thus, by (19), if the connected component C of $|[X/Q](\mathbb{R})|$ is an interval, then $g^{-1}(C)$ is an union of a intervals and b circles with $a + 2b \leq \#K$. Hence, we have:

$$h^*(g^{-1}(C)) \leq \#K(\mathbb{C}) \quad \text{if } C \text{ is an interval.} \quad (21)$$

Therefore, we have:

$$\begin{aligned}
h^*([X/H](\mathbb{R})) &= \sum_{C \in \pi_0([X/Q](\mathbb{R}))} h^*(g^{-1}(C)) \\
&= \sum_{C \text{ circle}} h^*(g^{-1}(C)) + \sum_{C \text{ interval}} h^*(g^{-1}(C)) \\
&\stackrel{(a)}{\leq} \sum_{C \text{ circle}} 2 \cdot \#K(\mathbb{C}) + \sum_{C \text{ interval}} \#K(\mathbb{C}) \\
&= \#K(\mathbb{C}) \cdot h^*([X/Q](\mathbb{R})) \\
&\stackrel{(b)}{\leq} \#K(\mathbb{C}) \cdot h^*([X/Q](\mathbb{C})) \\
&\stackrel{(c)}{=} h^*([X/H](\mathbb{C})),
\end{aligned}$$

where (a) holds by (20) and (21), (b) by the assumption that the Smith–Thom inequality (3) holds for $[X/Q]$, while (c) holds by Proposition 4.2, which we can apply since H is abelian. This proves Step 1. \square

Let $\Delta \subset X(\mathbb{C})/Q$ be the branch locus of the quotient map $q: X(\mathbb{C}) \rightarrow X(\mathbb{C})/Q$. For each $x \in \Delta$ choose an element $y_x \in X(\mathbb{C})$ such that $q(y_x) = x$. Define

$$\Delta' := \{x \in \Delta \cap (X/Q)(\mathbb{R}) \mid h^1(G, x) > 1\},$$

where $h^1(G, x) = \#H^1(G, H_{y_x}(\mathbb{C}))$ for some $y \in q^{-1}(x)$.

Step 2: *The Smith–Thom inequality (3) holds when the action of H on X over \mathbb{R} is faithful.*

Proof. Assume that H acts faithfully on X . By Lemma 4.8, the natural map $f: |\mathcal{X}(\mathbb{R})| \rightarrow M(\mathbb{R})$ is surjective. Let $C \subset (X/Q)(\mathbb{R})$ be a connected component which is homeomorphic to a circle. Since the action of H is faithful, by Proposition 4.9, $f^{-1}(C)$ is homeomorphic to a circle if $h^1(G, x) = 1$ for each $x \in C$, and $f^{-1}(C)$ is homeomorphic to the union of $\#(C \cap \Delta')$ intervals if $\Delta' \cap C \neq \emptyset$. In particular, we have:

$$h^*(f^{-1}(C)) = \begin{cases} 2 & \text{if } C \cap \Delta' = \emptyset, \\ \#(C \cap \Delta') & \text{if } C \cap \Delta' \neq \emptyset. \end{cases}$$

Let $I \subset (X/Q)(\mathbb{R})$ be a connected component which is homeomorphic to the open interval $(0, 1)$. Since the action of H is faithful, by Proposition 4.9, $f^{-1}(I)$ is homeomorphic

to the union of $\#(I \cap \Delta') + 1$ intervals. In particular, we have:

$$h^*(f^{-1}(I)) = \#(I \cap \Delta') + 1.$$

Therefore, we have:

$$\begin{aligned}
h^*(|\mathcal{X}(\mathbb{R})|) &= \sum_{C \in \pi_0(M(\mathbb{R})) \text{ circle}} h^*(f^{-1}(C)) + \sum_{I \in \pi_0(M(\mathbb{R})) \text{ interval}} h^*(f^{-1}(I)) \\
&\stackrel{(a)}{=} \sum_{C \cap \Delta' = \emptyset} 2 + \sum_{C \cap \Delta' \neq \emptyset} \#(C \cap \Delta') + \sum_I (\#(C \cap \Delta') + 1) \\
&= \left(\sum_{C \cap \Delta' = \emptyset} 2 + \sum_I 1 \right) + \left(\sum_{C \cap \Delta' \neq \emptyset} \#(C \cap \Delta') + \sum_{I \cap \Delta' \neq \emptyset} \#(C \cap \Delta') \right) \\
&\leq h^*((X/Q)(\mathbb{R})) + \sum_{x \in \Delta} 1 \\
&\stackrel{(b)}{\leq} h^*((X/Q)(\mathbb{R})) + \sum_{x \in \Delta} (\#(H_{y_x}(\mathbb{C})/H_{y_x}(\mathbb{C})) - 1) \\
&\stackrel{(c)}{\leq} h^*((X/Q)(\mathbb{C})) + \sum_{x \in \Delta} (\#(H_{y_x}(\mathbb{C})/H_{y_x}(\mathbb{C})) - 1) \\
&\stackrel{(d)}{=} h^*(|\mathcal{I}_{\mathcal{X}}(\mathbb{C})|).
\end{aligned}$$

Here, (a) follows from the previous discussion, (b) from the fact that we have $2 \leq \#(H_{y_x}(\mathbb{C})/H_{y_x}(\mathbb{C}))$, (c) from the classical Smith–Thom inequality (1) for X/Q , and, finally, (d) from Proposition 4.2. This proves Step 2. \square

By combining Steps 1 and 2, Theorem 1.4 follows. \square

5 Topology of split gerbes over a real variety

Let U be a geometrically connected real variety with $U(\mathbb{R}) \neq \emptyset$, and let $H \rightarrow U$ be a finite étale group scheme, equipped with the trivial action on U over U . In this section, we describe how to compute the topology of $|[U/H](\mathbb{R})|$ by comparing it with $U(\mathbb{R})$. This analysis leads to the proof of Theorem 1.5 as well as Corollaries 1.6 and 1.7.

In Section 5.1, we introduce notation that will be used throughout Section 5. The main work toward proving Theorem 1.5 is carried out in Section 5.2. In Section 5.3, we assemble all the ingredients and prove Theorem 1.5 along with Corollaries 1.6 and 1.7.

5.1 Notation and description of the complex inertia. Let U be a geometrically connected real variety with $U(\mathbb{R}) \neq \emptyset$, and let $H \rightarrow U$ be a finite étale group scheme, equipped with the trivial action on U over U . For $x \in U(\mathbb{R})$, we let H_x denote the fiber of $H \rightarrow U$ over x ; thus H_x is a finite étale group scheme over \mathbb{R} . We let $\bar{x}: \text{Spec}(\mathbb{C}) \rightarrow U$ be the geometric point associated to x , and define $H_{\bar{x}} = H \times_U \bar{x}$. Thus, $H_{\bar{x}}$ is the constant group scheme over \mathbb{C} associated to the finite group $H_x(\mathbb{C})$ which, by abuse of notation, we will also denote by $H_{\bar{x}}$. The finite group $H_{\bar{x}}$ is endowed with an action of G whose associated involution we denote by

$$\sigma_x: H_{\bar{x}} \rightarrow H_{\bar{x}}. \quad (22)$$

The natural map $U \rightarrow \mathcal{X}$ is a section of the coarse moduli space map $[U/H] \rightarrow U/H = U$, so that the map $f: |[U/H](\mathbb{R})| \rightarrow U(\mathbb{R})$ is surjective. Proposition 2.2 implies the following description of the complex locus of the inertia stack $\mathcal{I}_{[U/H]}$ of $[U/H]$.

Lemma 5.1. *In the above notation, we have*

$$|\mathcal{I}_{[U/H]}(\mathbb{C})| \simeq H(\mathbb{C})/H(\mathbb{C}) := \{(p, h) \in U(\mathbb{C}) \times H(\mathbb{C}) \mid h \in H_p(\mathbb{C})\} / \sim$$

where \sim denotes the equivalence relation $(p, h) \sim (p', h')$ if $p = p'$ and h is conjugate to h' in $H_p(\mathbb{C})$. In particular, when H is abelian, one has $|\mathcal{I}_{[U/H]}(\mathbb{C})| \simeq H(\mathbb{C})$. \square

5.2 Topological and algebraic fundamental groups

5.2.1 Action of fundamental groups. Fix $p \in U(\mathbb{R})$, and write C for the connected component of $U(\mathbb{R})$ containing p . The finite étale group scheme $H \rightarrow U$ corresponds to an action $\pi_1^{\text{ét}}(U, \bar{p})$ on $H_{\bar{p}}$,

$$\rho_p: \pi_1^{\text{ét}}(U, \bar{p}) \rightarrow \text{Aut}(H_{\bar{p}}),$$

which is compatible with the group structure of $H_{\bar{p}}$. Here, $\pi_1^{\text{ét}}(U, \bar{p})$ is the étale fundamental group of U at the geometric point \bar{p} .

Since U is geometrically connected and $G = \pi_1^{\text{ét}}(\text{Spec}(\mathbb{R}))$, the natural morphisms $U_{\mathbb{C}} \rightarrow U \rightarrow \text{Spec}(\mathbb{R})$ induce a short exact sequence of groups

$$1 \rightarrow \pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p}) \rightarrow \pi_1^{\text{ét}}(U, \bar{p}) \rightarrow G \rightarrow 1. \quad (23)$$

Restricting ρ_p to $\pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p})$, we get an action of $\pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p})$ on $H_{\bar{p}}$,

$$\rho_p^{\mathbb{C}}: \pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p}) \rightarrow \text{Aut}(H_{\bar{p}}),$$

which corresponds to the étale $U_{\mathbb{C}}$ -group scheme $H_{\mathbb{C}} \rightarrow U_{\mathbb{C}}$. Recall that $\pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p})$ identifies with the profinite completion of usual fundamental group $\pi_1(U(\mathbb{C}), p)$ so that, in particular, there is a natural map $\pi_1(U(\mathbb{C}), p) \rightarrow \pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p})$. We denote again by

$$\rho_p^{\mathbb{C}}: \pi_1(U(\mathbb{C}), p) \rightarrow \text{Aut}(H_{\bar{p}})$$

the restriction of $\rho_p^{\mathbb{C}}: \pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p}) \rightarrow \text{Aut}(H_{\bar{p}})$ along the map $\pi_1(U(\mathbb{C}), p) \rightarrow \pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p})$; this representation of $\pi_1(U(\mathbb{C}), p)$ corresponds to the topological covering $H(\mathbb{C}) \rightarrow U(\mathbb{C})$.

Viewing p as a morphism of schemes $p: \text{Spec}(\mathbb{R}) \rightarrow U$, we get a morphism $\pi_1(p): G = \pi_1^{\text{ét}}(\text{Spec}(\mathbb{R})) \rightarrow \pi_1^{\text{ét}}(U, \bar{p})$ which splits (23), and hence yields an isomorphism

$$\pi_1^{\text{ét}}(U, \bar{p}) \simeq \pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p}) \rtimes G. \quad (24)$$

Hence, viewing G as a subgroup $G \subset \pi_1^{\text{ét}}(U, \bar{p})$ via the isomorphism (24), this induces an action of $G = \langle \sigma \rangle$ on $\pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p})$ by the usual formula $\sigma \cdot \alpha = \sigma \alpha \sigma^{-1}$ for $\alpha \in \pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p})$. This action is compatible with the natural action of G on $\pi_1(U(\mathbb{C}), p)$ defined as follows: for $\alpha \in \pi_1(U(\mathbb{C}), p)$, we have $\sigma \cdot \alpha = \sigma_{U*}(\alpha)$.

Restricting ρ_p to G via $\pi_1(p)$, we get an action of G on $H_{\bar{p}}$ which corresponds to the natural involution σ_p on $H_{\bar{p}}$, see (22). Consider the morphism $\pi_1(C, p) \rightarrow \pi_1(U(\mathbb{C}), p)$ induced by the embedding $C \subset U(\mathbb{C})$, and define

$$\rho_p^C: \pi_1(C, p) \rightarrow \text{Aut}(H_{\bar{p}}) \quad (25)$$

as the composition of $\rho_p^{\mathbb{C}}$ with $\pi_1(C, p) \rightarrow \pi_1(U(\mathbb{C}), p)$.

Lemma 5.2. *The above action (25) of $\pi_1(C, p)$ on $H_{\bar{p}}$ commutes with σ_p , in the sense that $\sigma_p(\gamma \cdot x) = \gamma \cdot \sigma_p(x)$ for $\gamma \in \pi_1(C, p)$ and $x \in H_{\bar{p}}$. In particular, it preserves the subset $Z^1(G, H_{\bar{p}}) = \{x \in H_{\bar{p}} \mid x \cdot \sigma_p(x) = e\} \subset H_{\bar{p}}$. Moreover, the induced action of $\pi_1(C, p)$ on $Z^1(G, H_{\bar{p}})$ descends to an action of $\pi_1(C, p)$ on $H^1(G, H_{\bar{p}})$.*

Proof. We need to show that for every $\alpha \in \pi_1(C, p)$, one has

$$\rho_p^C(\alpha) \circ \sigma_p = \sigma_p \circ \rho_p^C(\alpha) \quad \text{as maps} \quad H_{\bar{p}} \rightarrow H_{\bar{p}}. \quad (26)$$

Via the isomorphism (24), we write each element $\beta \in \pi_1^{\text{ét}}(U, \bar{p})$ as a pair $\beta = (\beta_1, \beta_2)$ with $\beta_1 \in \pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p})$ and $\beta_2 \in G$. Denoting again by α the image of α in $\pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p})$, equation (26) can be rewritten as

$$\rho_p(\alpha, e)^{-1} \circ \rho_p(e, \sigma) \circ \rho_p(\alpha, e) = \rho_p(e, \sigma),$$

where e is the neutral element of $\pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p})$. Since ρ_p is a group homomorphism, we have

$$\rho_p(\alpha, e)^{-1} \circ \rho_p(e, \sigma) \circ \rho_p(\alpha, e) = \rho_p((\alpha^{-1}, e) \cdot (e, \sigma) \cdot (\alpha, e)).$$

By the definition of the semi-direct product group structure, we have

$$(\alpha^{-1}, e) \cdot (e, \sigma) \cdot (\alpha, e) = (\alpha^{-1} \sigma^{-1} \alpha \sigma, \sigma).$$

The image $\alpha \in \pi_1(U(\mathbb{C}), p)$ of $\alpha \in \pi_1(C, p)$ satisfies $\sigma_{U*}(\alpha) = \sigma_U \circ \alpha = \alpha$. For the image $\alpha \in \pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p})$, one therefore has $\sigma \cdot \alpha = \sigma^{-1} \alpha \sigma = \alpha$. Hence, we get

$$\rho_p(\alpha, e)^{-1} \circ \rho_p(e, \sigma) \circ \rho_p(\alpha, e) = \rho_p((\alpha^{-1}, e)(e, \sigma)(\alpha, e)) = \rho_p(e, \sigma),$$

and the proof is concluded. \square

5.2.2 Change of base point. In the previous section, we fixed a point $p \in U(\mathbb{R})$ to study the fiber H_p . For concrete calculations, it will be important to understand how H_p varies when the point p varies in $U(\mathbb{R})$. This is the content of the following proposition.

Proposition 5.3. *Let U be a geometrically connected scheme of finite type over \mathbb{R} with $U(\mathbb{R}) \neq \emptyset$. Let $Y \rightarrow U$ be a finite étale cover. Let $p, q \in U(\mathbb{R})$ and choose a topological path $\gamma_{q,p}$ from q to p in $U(\mathbb{C})$. Define $\omega_{q,p} := (\gamma_{q,p} \circ \sigma_U) * \gamma_{q,p}^{-1} \in \pi_1(U(\mathbb{C}), p)$ (where $*$ denotes the composition of paths). Then the following diagram commutes:*

$$\begin{array}{ccc} Y_{\bar{q}} & \xrightarrow{(\gamma_{q,p})_*} & Y_{\bar{p}} \\ \downarrow \sigma_q & & \downarrow \sigma_p \\ Y_{\bar{q}} & & Y_{\bar{p}} \\ \parallel & & \downarrow \omega_{q,p} \\ Y_{\bar{q}} & \xrightarrow{(\gamma_{q,p})_*} & Y_{\bar{p}} \end{array}$$

Here, $\omega_{q,p}$ acts on $Y_{\bar{p}}$ as an element of $\pi_1(U(\mathbb{C}), p)$, and $(\gamma_{q,p})_*: Y_{\bar{q}} \xrightarrow{\sim} Y_{\bar{p}}$ is the canonical isomorphism induced by the path $\gamma_{q,p}$.

Proof. For any path $\gamma_{q,p}: [0, 1] \rightarrow U(\mathbb{C})$ from q to p , and any point $y \in Y_{\bar{q}}$, we let $\tilde{\gamma}_{q,p}^y$ be the unique path in $Y(\mathbb{C})$ that lifts $\gamma_{q,p}$ and that satisfies $\tilde{\gamma}_{q,p}^y(0) = y$. This yields an isomorphism

$$(\gamma_{q,p})_*: Y_{\bar{q}} \xrightarrow{\sim} Y_{\bar{p}}, \quad y \mapsto \tilde{\gamma}_{q,p}^y(1).$$

By construction, we have:

$$\sigma_Y((\gamma_{q,p})_*(y)) = \sigma_Y(\tilde{\gamma}_{q,p}^y(1)) \quad \text{and} \quad \omega_{q,p} \cdot (\gamma_{q,p})_*(\sigma_Y(y)) = \omega_{q,p} \cdot \tilde{\gamma}_{q,p}^{\sigma_Y(y)}.$$

Observe that

$$\sigma_Y(\tilde{\gamma}_{q,p}^y(1)) = (\sigma_{U*}(\tilde{\gamma}_{p,q}^y))(1)$$

and that $\sigma_{U*}(\tilde{\gamma}_{p,q}^y)$ is a path in $Y(\mathbb{C})$ that lifts $\sigma_{U*}(\gamma_{q,p})$ and that starts at $\sigma_Y(y)$. In other words,

$$\sigma_Y((\gamma_{q,p})_*(y)) = \widetilde{\sigma_{U*}(\gamma_{q,p})}^{\sigma_Y(y)}(1).$$

On the other hand, by construction of the action of $\pi_1(U(\mathbb{C}), p)$ on $Y_{\bar{p}}$, one has

$$\omega_{q,p} \cdot \left(\widetilde{\gamma_{q,p}^{\sigma_Y(y)}}(1) \right) = \widetilde{\omega_{q,p} * \gamma_{q,p}}^{\sigma_Y(y)}(1).$$

But $\omega_{q,p} * \gamma_{q,p} = \sigma_{U*}(\gamma_{q,p})$ by definition of $\omega_{q,p}$, hence the proof is concluded. \square

Example 5.4. Let $U \subset \mathbb{G}_m$ be an open subset whose real part contains $[-1, 0)$ and $(0, 1]$ and let $\pi: E \rightarrow U$ be a family of elliptic curves over \mathbb{R} . Let $p = 1 \in U(\mathbb{R})$ and assume that Y_p is a maximal real elliptic curve (so that $Y_p(\mathbb{R})$ has two connected components). Consider the local system $\pi_*\mathbb{Z}/2$ of finite dimensional $\mathbb{Z}/2$ -modules on $U_{\text{ét}}$, and let

$$Y \rightarrow U$$

be the associated finite étale cover. Thus,

$$Y_{\bar{q}} = H^1(E_q(\mathbb{C}), \mathbb{Z}/2) \quad \text{for} \quad q \in U(\mathbb{R}).$$

Since E_p is a maximal real elliptic curve, the action of G on $Y_{\bar{p}} = H^1(E_p(\mathbb{C}), \mathbb{Z}/2)$ is trivial.

1. Assume that the action of the standard loop γ around 0 (viewed as an element of $\pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p})$) on $H^1(E_p(\mathbb{C}), \mathbb{Z}/2)$ is not trivial (this happens for example for the family whose affine equation is $y^2 = (x^2 - t)(x + 2)$ where t is the coordinate of U). Let $q = -1$ and choose as $\gamma_{q,p}$ the standard "half circle" around 0, so

that $\omega_{q,p} = \gamma$, hence it acts non-trivially on $\pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p})$. Since the action of G on $H^1(E_p(\mathbb{C}), \mathbb{Z}/2)$ is trivial, we deduce from Proposition 5.3 that the action of G on $H^1(E_q(\mathbb{C}), \mathbb{Z}/2)$ is not trivial. In particular, the real elliptic curve E_q is not maximal.

2. Assume that the action of $\pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p})$ on $H^1(E_p(\mathbb{C}), \mathbb{Z}/2)$ is trivial (this happens for example for the family whose affine equation is $y^2 = x(x+t)(x+2)$ where t is the coordinate of U). Let $q = -1$. Since $\pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p})$ acts trivially on $H^1(E_{\bar{p}}(\mathbb{C}), \mathbb{Z}/2)$, for every choice of path $\gamma_{q,p}$ from -1 to 1 , the loop ω acts trivially on $H^1(E_p(\mathbb{C}), \mathbb{Z}/2)$, so that from Proposition 5.3, we deduce that E_q is maximal.

Corollary 5.5. *Let U be a geometrically connected scheme locally of finite type over \mathbb{R} with $U(\mathbb{R}) \neq \emptyset$, and let $H \rightarrow U$ be a finite étale group scheme. Let $C \subset U(\mathbb{R})$ be a connected component. For $q, p \in C$, let $\gamma_{q,p}: [0, 1] \rightarrow C$ be a path from q to p . The induced isomorphism $(\gamma_{q,p})_*: H_{\bar{q}} \simeq H_{\bar{p}}$ is an isomorphism of finite G -groups.*

Proof. Define $\omega_{q,p}$ as in Proposition 5.3. Then

$$\omega_{q,p} = (\gamma_{q,p} \circ \sigma_U) * \gamma_{q,p}^{-1} = \gamma_{q,p} * \gamma_{q,p}^{-1} = e \in \pi_1(U(\mathbb{C}), p).$$

Thus, the corollary follows from Proposition 5.3. \square

5.3 Proof of Theorem 1.5 and its corollaries. We are ready to proceed with the:

Proof of Theorem 1.5. Item 1 of the theorem is the combination of [AGF25, Theorem 1.5] and Proposition 2.1 (alternatively, see [AGF25, Lemma 5.8 & Proposition 5.9]). Item 2 is the content of Lemma 5.2. Finally, item 3 follows from the previous items and from [AGF25, Lemma 5.8 & Proposition 5.9], and we are done. \square

Proof of Corollary 1.6. By Lemma 5.1, $|\mathcal{I}_{\mathcal{X}}(\mathbb{C})|$ is homeomorphic to $H(\mathbb{C})$, so we need to prove that

$$h^*(|[U/H](\mathbb{R})|) \leq h^*(H(\mathbb{C})),$$

Consider the diagram of finite étale covers

$$\begin{array}{ccc} & & H(\mathbb{R}) \\ & & \downarrow g \\ |[U/H](\mathbb{R})| & \xrightarrow{f} & U(\mathbb{R}) \end{array}$$

We claim that there exists a canonical embedding of sheaves

$$f_*\mathbb{Z}/2 \hookrightarrow g_*\mathbb{Z}/2. \quad (27)$$

To prove this, we may fix a connected component $C \subset U(\mathbb{R})$, and define the embedding after restricting both sheaves to C . Fix $p \in C$. By assumption, we have an isomorphism of G -groups $H_{\bar{p}} = A$ for a finite G -module A with underlying abelian group $(\mathbb{Z}/2)^n$. Let $\sigma: A \rightarrow A$ be the involution corresponding to the G -module structure of A .

On the one hand, the étale cover $g: H(\mathbb{R})|_C \rightarrow C$ has fiber $g^{-1}(p) = A^\sigma$ and corresponds to the action of $\pi_1(C, p)$ on A^σ induced by the map $\pi_1(C, p) \rightarrow \pi_1(U(\mathbb{C}), p)$ and the action of $\pi_1(U(\mathbb{C}), p)$ on A (which is, in turn, induced by the étale cover $H \rightarrow U$). On the other hand, the étale cover $f^{-1}(C) \rightarrow C$ corresponds by Theorem 1.5 to the action of $\pi_1(C, p)$ on $f^{-1}(p) = H^1(G, A)$ induced by the morphism $\pi_1(C, p) \rightarrow \pi_1(U(\mathbb{C}), p)$ and the action of $\pi_1(U(\mathbb{C}), p)$ on A .

Note that, since A is an \mathbb{F}_2 -vector space, one has

$$H^1(G, A) = A^\sigma / (1 + \sigma)A.$$

Moreover, by the above, the quotient map

$$A^\sigma \rightarrow A^\sigma / (1 + \sigma)A = H^1(G, A) \quad (28)$$

is $\pi_1(C, p)$ -equivariant. Applying the contravariant functor $(\mathbb{Z}/2)^{(-)}$ to (28) gives a $\pi_1(C, p)$ -equivariant embedding

$$(f_*\mathbb{Z}/2)_p = (\mathbb{Z}/2)^{H^1(G, A)} \hookrightarrow (\mathbb{Z}/2)^{A^\sigma} = (g_*\mathbb{Z}/2)_p.$$

Since $f_*\mathbb{Z}/2$ and $g_*\mathbb{Z}/2$ are locally constant sheaves, this yields the desired embedding of sheaves (27).

By taking global sections of the embedding of sheaves (27), we obtain the inequality

$$\dim H^0([U/H](\mathbb{R}), \mathbb{Z}/2) \leq \dim H^0(H(\mathbb{R}), \mathbb{Z}/2). \quad (29)$$

Since f and g are finite maps, we have $f_* = f_!$ and $g_* = g_!$. Hence, we can take global sections with compact support, to get

$$\dim H_c^0([U/H](\mathbb{R}), \mathbb{Z}/2) \leq \dim H_c^0(H(\mathbb{R}), \mathbb{Z}/2). \quad (30)$$

All the connected components of $U(\mathbb{R})$ are open intervals or circles; as $f: |[U/H](\mathbb{R})| \rightarrow U(\mathbb{R})$ is a finite topological covering by Theorem 1.5, this also holds for $|[U/H](\mathbb{R})|$. Hence, one has:

$$\begin{aligned} h^*([U/H](\mathbb{R})) &= \dim H^0([U/H](\mathbb{R}), \mathbb{Z}/2) + \dim H_c^0([U/H](\mathbb{R}), \mathbb{Z}/2) \\ &\leq \dim H^0(H(\mathbb{R}), \mathbb{Z}/2) + \dim H_c^0(H(\mathbb{R}), \mathbb{Z}/2) \\ &= h^*(H(\mathbb{R})), \end{aligned}$$

where the inequality follows from (29) and (30). The classical Smith–Thom inequality (1) implies that $h^*(H(\mathbb{R})) \leq h^*(H(\mathbb{C}))$, so that

$$h^*([U/H](\mathbb{R})) \leq h^*(H(\mathbb{R})) \leq h^*(H(\mathbb{C})),$$

and this concludes the proof. \square

Proof of Corollary 1.7. Since U is a smooth proper curve, by Theorem 1.5, $|[U/H](\mathbb{R})|$ is a disjoint union of circles, so that

$$h^*([U/H](\mathbb{R})) = 2 \cdot \#\pi_0([U/H](\mathbb{R})).$$

By Theorem 1.5, for every connected component $C_i \in \pi_0([U/H](\mathbb{R}))$, the map $f^{-1}(C_i) \rightarrow C_i$ is the étale cover corresponding to the action of $\pi_1(C_i, p_i)$ on $H^1(G, H_{\bar{p}_i})$. Hence, $f^{-1}(C_i)$ has $\#(H^1(G, H_{\bar{p}_i})/\pi_1(C_i, p_i))$ connected components. Consequently, we get

$$h^*([U/H](\mathbb{R})) = \sum_{i=1}^m 2 \cdot \# \left(\frac{H^1(G, H_{\bar{p}_i})}{\pi_1(C_i, p_i)} \right). \quad (31)$$

Assuming Conjecture 1.1, we have

$$h^*([U/H](\mathbb{R})) \leq h^*(I_{[U/H]}(\mathbb{C})). \quad (32)$$

Therefore, to finish the proof, it suffices to show that

$$\begin{aligned} h^*(I_{[U/H]}(\mathbb{C})) &= 2 \cdot \# \left(\frac{H_{\bar{p}}}{\pi_1(U(\mathbb{C}), p)} \right) \\ &\quad + \sum_{D \in (H_{\bar{p}}/\pi_1(U(\mathbb{C}), p))} 2 \cdot (\#D \cdot (g-1) + 1), \end{aligned} \quad (33)$$

for indeed, by the fact that $\sum_D \#D = \#H_{\bar{p}}$, where the sum ranges over the orbits

$D \in (H_{\bar{p}}/\pi_1(U(\mathbb{C}), p))$, equations (31), (32) and (33) together imply equation (4).

By Lemma 5.1, the map $|I_{[U/H]}(\mathbb{C})| = H(\mathbb{C}) \rightarrow |[U/H](\mathbb{C})| = U(\mathbb{C})$ is the étale cover corresponding to the action of $\pi_1(U(\mathbb{C}), p)$ on $H_p(\mathbb{C})$. Hence, $|I_{[U/H]}(\mathbb{C})|$ has $\#(H_{\bar{p}}/\pi_1(U(\mathbb{C}), p))$ connected components, and each of these is a connected étale cover whose degree equals the cardinality of the orbit. Thus, (33) follows from the Riemann–Hurwitz formula, and the proof is finished. \square

6 Gerbes and the homotopy exact sequence

In this section, we refine the results of Section 5, in order to be able to apply them in explicit examples. A key step is the reinterpretation of Proposition 5.3 in terms of splittings of the homotopy exact sequence (23). This reinterpretation is developed in Section 6.1 and subsequently applied in Section 6.2 to the study of certain gerbes over an Enriques surface.

We retain the notation introduced in Section 5. In particular, U denotes a geometrically connected scheme locally of finite type over \mathbb{R} with $U(\mathbb{R}) \neq \emptyset$, and $H \rightarrow U$ is a finite étale group scheme over U .

6.1 Topological paths and splitting of the homotopy exact sequence

6.1.1 Galois formalism. Write \mathbf{Fset} for the category of finite sets and, for a scheme Z , $\mathbf{Fét}(Z)$ for the category of finite étale covers of Z . Recall that for every $q \in U(\mathbb{R})$, the group $\pi_1^{\text{ét}}(U, \bar{q})$ (resp. $\pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{q})$) is the automorphism of the functor

$$(-)_{\bar{q}} : \mathbf{Fét}(U) \rightarrow \mathbf{Fset} \quad (\text{resp. } (-)_{\bar{q}}^{\mathbb{C}} : \mathbf{Fét}(U_{\mathbb{C}}) \rightarrow \mathbf{Fset})$$

sending $Y \rightarrow U$ (resp. $Y \rightarrow U_{\mathbb{C}}$) to the geometric fiber $Y_{\bar{q}}$. By the general formalism of Galois categories, every isomorphism of functors

$$\phi : (-)_{\bar{q}} \xrightarrow{\cong} (-)_{\bar{p}},$$

induces an isomorphism, well defined up to conjugation,

$$\varphi : \pi_1^{\text{ét}}(U, \bar{q}) \xrightarrow{\cong} \pi_1^{\text{ét}}(U, \bar{p}),$$

in such a way that the action of G on $H_{\bar{q}}$ is induced by the action of $\pi_1^{\text{ét}}(U, \bar{p})$ on $H_{\bar{p}}$ and the composition

$$G = \pi_1^{\text{ét}}(\text{Spec}(\mathbb{R})) \xrightarrow{\pi_1(q)} \pi_1^{\text{ét}}(U, \bar{q}) \xrightarrow{\varphi} \pi_1^{\text{ét}}(U, \bar{p}).$$

Remark 6.1. Let us emphasize that, in general, the isomorphism $\varphi: \pi_1^{\text{ét}}(U, \bar{q}) \xrightarrow{\sim} \pi_1^{\text{ét}}(U, \bar{p})$ induced by an isomorphism of functors $\phi: (-)_{\bar{q}} \xrightarrow{\sim} (-)_{\bar{p}}$ is not unique. It is only unique up to conjugation.

Since both $\pi_1(p)$ and $\varphi \circ \pi_1(q)$ are splittings of the exact sequence

$$0 \rightarrow \pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p}) \rightarrow \pi_1^{\text{ét}}(U, \bar{p}) \rightarrow G \rightarrow 0,$$

to understand the action of G on $H_{\bar{q}}$, one has to understand how the different splittings of this sequence are related. This is the main result of the section.

6.1.2 Splittings of the homotopy exact sequence. Let $p, q \in U(\mathbb{R})$. Let

$$\gamma_{q,p}: [0, 1] \rightarrow U(\mathbb{C})$$

be a path from q to p . The isomorphisms

$$(\gamma_{q,p})_*: Y_{\bar{q}} \simeq Y_{\bar{p}}$$

induced by $\gamma_{q,p}$ fit together to give an isomorphism $\varphi_{q,p}^{\mathbb{C}}: (-)_{\bar{q}}^{\mathbb{C}} \xrightarrow{\sim} (-)_{\bar{p}}^{\mathbb{C}}$ of fiber functors. In turn, this induces an isomorphism

$$\varphi_{q,p}^{\mathbb{C}}: \pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{q}) \xrightarrow{\sim} \pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p}),$$

well defined up to conjugation, extending the usual isomorphism $\pi_1(U(\mathbb{C}), q) \rightarrow \pi_1(U(\mathbb{C}), p)$ defined by $\alpha \mapsto \gamma_{q,p} \alpha \gamma_{q,p}^{-1}$. Write

$$\omega := \sigma_{U*}(\gamma_{q,p}) * \gamma_{q,p}^{-1} \in \pi_1(U(\mathbb{C}), p),$$

where $*$ is the concatenation of paths. By abuse of notation, let $\omega \in \pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p})$ be the image of ω under the natural morphism $\pi_1(U(\mathbb{C}), p) \rightarrow \pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p})$.

Proposition 6.2. *In the above notation, consider the map $f: G \rightarrow \pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p}) \rtimes G$*

defined as the composition

$$f: G \xrightarrow{\pi_1(q)} \pi_1^{\text{ét}}(U, \bar{q}) \xrightarrow{\varphi_{q,p}} \pi_1^{\text{ét}}(U, \bar{p}) \simeq \pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p}) \rtimes G,$$

where the isomorphism on the right is defined by the splitting of the homotopy exact sequence (23) induced by the section $\pi_1(p): G \rightarrow \pi_1^{\text{ét}}(U, \bar{p})$. Then $f(\sigma)$ is conjugate to (ω, σ) .

Proof. Let $Y \rightarrow U$ be a finite connected étale cover. By Proposition 5.3, the action of $G = \pi_1^{\text{ét}}(\text{Spec}(\mathbb{R}))$ on $Y_{\bar{p}}$ induced by the action of $\pi_1^{\text{ét}}(U, \bar{p})$ on $Y_{\bar{p}}$, and up to multiplying by ω , the composition

$$G \xrightarrow{\pi_1(q)} \pi_1^{\text{ét}}(U, \bar{q}) \xrightarrow{\varphi_{q,p}} \pi_1^{\text{ét}}(U, \bar{p})$$

identifies with the natural action of G on $Y_{\bar{p}}$. To be precise, after identifying $Y_{\bar{p}}$ and $Y_{\bar{q}}$ using $\gamma_{q,p}$, one has $\sigma_q = \omega \cdot \sigma_p$. The proposition follows from this. \square

Examples 6.3. Let $U = \mathbb{G}_m$ and take $p = 1 \in \mathbb{G}_m(\mathbb{R})$. In this case, the étale fundamental group is given by

$$\pi_1^{\text{ét}}(U, \bar{p}) \simeq \hat{\mathbb{Z}} \rtimes G,$$

where G acts on $\hat{\mathbb{Z}}$ by inversion.

1. Take $q = 2$ and choose $\gamma_{q,p}$ as the natural path contained in the real part connecting q to p . In this case, ω is the trivial loop, so the image of the section corresponding to $\pi_1(q)$ is identified with $(0, e)$.
2. Again, take $q = 2$, but this time let $\gamma_{q,p}$ be a loop not contained in the real part, such that ω is nontrivial (for example, the path shown on the left in Figure 5). By construction, the class of ω in $\pi_1(U(\mathbb{C}), p) \simeq \mathbb{Z}$ is 2, so under this choice of γ , the image of the section $\pi_1(q)$ is $(2, e)$. Although different from the previous case, we note that $(2, e)$ is conjugate to $(0, e)$ in $\mathbb{Z} \rtimes G$, meaning that the conjugacy class of the section remains unchanged (see Remark 6.1).
3. Take $q = -1$ and choose $\gamma_{q,p}$ as the "half-circle path" from -1 to 1 , as depicted on the right in Figure 5. In this case, γ corresponds to the class of 1 in \mathbb{Z} , so the image of the section $\pi_1(q)$ is $(1, e)$. Since $(1, e)$ is not conjugate to $(0, e)$, this section is genuinely different from the previous ones, even up to conjugation.




Figure 5: Two paths in $\mathbb{G}_m(\mathbb{C})$.

6.2 Smith–Thom for various split gerbes over an Enriques surface. In this section, we apply the theory developed in the previous section – particularly Theorem 1.5 – to verify the Smith–Thom inequality for certain gerbes over an Enriques surface.

Let U be an Enriques surface such that $U(\mathbb{R}) \neq \emptyset$, so that its K3 cover $h: V \rightarrow U$ is defined over \mathbb{R} . To simplify the discussion, we also assume that $V(\mathbb{R}) \neq \emptyset$. Fix a point $p \in U(\mathbb{R})$ in the image of $h: V(\mathbb{R}) \rightarrow U(\mathbb{R})$. Recall that $\pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p}) \simeq \mathbb{Z}/2$, so that, since $\mathbb{Z}/2$ has no non-trivial automorphisms, the section $\pi_1(p)$ induces an isomorphism

$$\pi_1^{\text{ét}}(U, \bar{p}) \simeq \pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p}) \times G,$$

such that the K3 cover $h: V \rightarrow U$ corresponds to the morphism

$$\pi_1: \pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p}) \times G \rightarrow \mathbb{Z}/2$$

which is the composition of the projection onto the first factor and the isomorphism $\pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p}) \simeq \mathbb{Z}/2$.

We let 1 be the generator of $\pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p})$. For every $q \in U(\mathbb{R})$, the group G acts trivially on $V_{\bar{q}}$ if and only if q is in the image of $h: V(\mathbb{R}) \rightarrow U(\mathbb{R})$. Consequently, by Proposition 6.2, the element ϵ_q corresponding to the section associated with q is given by

$$\epsilon_q = \begin{cases} (0, 0) & \text{if } q \text{ is in the image of } h: V(\mathbb{R}) \rightarrow U(\mathbb{R}), \\ (1, 0) & \text{otherwise.} \end{cases}$$

For the remainder of this section, we let $G = \mathbb{Z}/2$ act on $\mathbb{Z}/2 \oplus \mathbb{Z}/2$ by exchanging the coordinates.

Example 6.4. Assume that $U(\mathbb{R})$ is the union of four copies of $\mathbb{P}^2(\mathbb{R})$ and two spheres

S^2 , and that the map $h: V(\mathbb{R}) \rightarrow U(\mathbb{R})$ is surjective (such Enriques surfaces exist, as shown in [DIK00, Table 8, page 180]).

Let $\pi_1: \pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p}) \times G \rightarrow \mathbb{Z}/2$ be the composition of the projection onto the first coordinate and the isomorphism $\pi_1^{\text{ét}}(U_{\mathbb{C}}, \bar{p}) \simeq \mathbb{Z}/2$, and let $\mathbb{Z}/2 \times G$ act on $\mathbb{Z}/2 \oplus \mathbb{Z}/2$ via this map. Denote by $H \rightarrow U$ the corresponding group scheme. Since $h: V(\mathbb{R}) \rightarrow U(\mathbb{R})$ is surjective, for every $q \in U(\mathbb{R})$, the image of the section $\pi_1(q)$ is $(0, e)$. Consequently, the action of G on $H_{\bar{q}}$ is trivial, implying that $H^1(G, H_{\bar{q}}) = \mathbb{Z}/2 \times \mathbb{Z}/2$. Let C be a connected component homeomorphic to S^2 . Since S^2 is simply connected, the cover $f^{-1}(C) \rightarrow C$ is trivial. Thus, the preimage of each such C under the map

$$f: |[U/H](\mathbb{R})| \rightarrow U(\mathbb{R})$$

consists of four copies of S^2 .

On the other hand, let C be a connected component homeomorphic to $\mathbb{P}^2(\mathbb{R})$. The natural map $\pi_1(C) \rightarrow \pi_1(U(\mathbb{C}))$ is an isomorphism, as there is at least one spherical connected component in the real locus of the K3 cover over C (see the discussion in [DIK00, Section 3.5]). Thus, the cover $f^{-1}(C) \rightarrow C$ has three connected components, corresponding to the orbits $\{(0, 0)\}, \{(1, e)\}, \{(0, e), (1, 0)\}$ of the action of $\pi_1(U(\mathbb{C}))$ on $H^1(G, H_{\bar{q}}) = \mathbb{Z}/2 \times \mathbb{Z}/2$. Since the $\pi_1(C)$ -action on $\{(0, e), (1, 0)\}$ is nontrivial, the corresponding cover is homeomorphic to the universal cover $S^2 \rightarrow \mathbb{P}^2(\mathbb{R})$. Hence, $f^{-1}(C)$ is homeomorphic to the disjoint union of two copies of $\mathbb{P}^2(\mathbb{R})$ and one S^2 .

In conclusion, $[U/H](\mathbb{R})$ is homeomorphic to the disjoint union of:

- four copies of $S^2 \amalg \mathbb{P}^2(\mathbb{R}) \amalg \mathbb{P}^2(\mathbb{R})$, each lying over a $\mathbb{P}^2(\mathbb{R})$, and
- two copies of $\amalg_{1 \leq i \leq 4} S^2$, each lying over an S^2 .

In particular, we obtain

$$h^*([U/H](\mathbb{R})) = 4 \cdot (2 + 3 + 3) + 2 \cdot 8 = 48.$$

By Lemma 5.1, the inertia $|\mathcal{I}_{[U/H]}(\mathbb{C})| \rightarrow U(\mathbb{C})$ corresponds to the cover associated with the action of $\pi_1(U(\mathbb{C}))$ on $H_{\bar{p}}$, which has three connected components, corresponding to the orbits $\{(0, 0)\}, \{(1, e)\}, \{(0, e), (1, 0)\}$. Hence, $|\mathcal{I}_{[U/H]}(\mathbb{C})|$ is the disjoint union of two copies of $U(\mathbb{C})$ and one copy of its K3 cover. In particular,

$$h^*(|\mathcal{I}_{[U/H]}(\mathbb{C})|) = 16 \cdot 2 + 24 = 56.$$

Thus, the Smith–Thom inequality (3) is verified for the stack $[U/H]$.

Example 6.5. Retaining the notation of Example 6.4, we now assume that the image of the map $V(\mathbb{R}) \rightarrow U(\mathbb{R})$ consists of only three copies of $\mathbb{P}^2(\mathbb{R})$ and a single S^2 (such Enriques surfaces exist by [DIK00, Table 8, page 180]).

The description of the cover $f^{-1}(C) \rightarrow C$ remains the same for the connected components in the image of $V(\mathbb{R}) \rightarrow U(\mathbb{R})$. However, it differs for the connected components C_1 and C_2 , which are respectively homeomorphic to S^2 and $\mathbb{P}^2(\mathbb{R})$ but are not in the image. In the end, $|[U/H](\mathbb{R})|$ has the following description.

- Three copies of $S^2 \amalg \mathbb{P}^2(\mathbb{R}) \amalg \mathbb{P}^2(\mathbb{R})$, each lying over a $\mathbb{P}^2(\mathbb{R})$;
- One copy of $\amalg_{1 \leq i \leq 4} S^2$, lying over an S^2 ;
- One copy of $\mathbb{P}^2(\mathbb{R})$, lying over an $\mathbb{P}^2(\mathbb{R})$ and one copy of S^2 , lying over an S^2 .

This is illustrated in Figure 6 below, where the dark disks represent copies of $\mathbb{P}^2(\mathbb{R})$, and the light spheres represent copies of S^2 .




Figure 6: The morphism $|[U/H](\mathbb{R})| \rightarrow U(\mathbb{R})$

To justify this, choose a point $q_i \in C_i$. Since q_i is not in the image of $V(\mathbb{R}) \rightarrow U(\mathbb{R})$, the action of G on $V_{\bar{q}_i}$ is nontrivial. Consequently, the image of the section $\pi_1(q_i)$ is $(1, e)$, implying that G acts on $H_{\bar{q}_i}$ by exchanging the coordinates. In particular, $H^1(G, H_{\bar{q}_i}) = 0$ is trivial, so that $f^{-1}(C_i) \rightarrow C_i$ is an isomorphism. As in the previous example, one verifies that the Smith–Thom inequality holds in this case as well.

References

- [AGF25] Emiliano Ambrosi and Olivier de Gaay Fortman. *Topological groupoids with involution and real algebraic stacks*. 2025.
- [Alp25] Jarod Alper. *Stacks and Moduli*. <https://sites.math.washington.edu/~jarod/moduli.pdf>. 2025.
- [BD17] Indranil Biswas and Shane D’Mello. “M-curves and symmetric products”. In: *Proc. Indian Acad. Sci. Math. Sci.* 127.4 (2017), pp. 615–624.
- [Bor60] Armand Borel. *Seminar on Transformation Groups*. Annals of Mathematics Studies, No. 46. With contributions by G. Bredon, E. E. Floyd, D. Montgomery, R. Palais. Princeton University Press, Princeton, NJ, 1960, pp. vii+245.
- [BS22] Erwan Brugallé and Florent Schaffhauser. “Maximality of moduli spaces of vector bundles on curves”. In: *Épjournal Géom. Algébrique* 6 (2022), Art. 24, 15.
- [DIK00] Alex Degtyarev, Ilia Itenberg, and Viatcheslav Kharlamov. *Real Enriques Surfaces*. Vol. 1746. Lecture Notes in Mathematics. Springer-Verlag, Berlin, 2000, pp. xvi+259.
- [Flo52] Edwin Earl Floyd. “On periodic maps and the Euler characteristics of associated spaces”. In: *Trans. Amer. Math. Soc.* 72 (1952), pp. 138–147.
- [Fra18] Matthias Franz. “Symmetric products of equivariantly formal spaces”. In: *Canad. Math. Bull.* 61.2 (2018), pp. 272–281.
- [Fu23] Lie Fu. *Maximal real varieties from moduli constructions*. To appear in *Moduli*. 2023. arXiv: 2303.03368 [math.AG].
- [GF22a] Olivier de Gaay Fortman. “Moduli spaces and algebraic cycles in real algebraic geometry”. PhD thesis. École normale supérieure de Paris, 2022.

- [GF22b] Olivier de Gaay Fortman. “Real moduli spaces and density of non-simple real abelian varieties”. In: *Q. J. Math.* 73.3 (2022), pp. 969–989.
- [KR24] Viatcheslav Kharlamov and Rareş Răşdeaconu. *Unexpected loss of maximality: the case of Hilbert square of real surfaces*. 2024. arXiv: 2303.02796 [math.AG].
- [Man17] Frédéric Mangolte. *Variétés Algébriques Réelles*. Vol. 24. Cours Spécialisés. Société Mathématique de France, 2017, pp. vii+484.
- [Ser94] Jean-Pierre Serre. *Cohomologie Galoisienne*. Fifth. Vol. 5. Lecture Notes in Mathematics. Springer-Verlag, Berlin, 1994, pp. x+181.
- [Sko01] Alexei Skorobogatov. *Torsors and Rational Points*. Vol. 144. Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 2001, pp. viii+187.
- [Tho65] René Thom. “Sur l’homologie des variétés algébriques réelles”. In: *Differential and Combinatorial Topology (A Symposium in Honor of Marston Morse)*. Princeton Univ. Press, Princeton, N.J., 1965, pp. 255–265.

EMILIANO AMBROSI, INSTITUT DE RECHERCHE MATHÉMATIQUE AVANCÉE (IRMA),
 7 RUE RENÉ DESCARTES, 67084 STRASBOURG, FRANCE
E-mail address: `eambrosi@unistra.fr`

OLIVIER DE GAAY FORTMAN, DEPARTMENT OF MATHEMATICS, UTRECHT UNIVERSITY,
 BUDAPESTLAAN 6, 3584 CD UTRECHT, THE NETHERLANDS
E-mail address: `a.o.d.degaayfortman@uu.nl`